

Bruce A. Measure
Chair
Montana

Rhonda Whiting
Montana

W. Bill Booth
Idaho

James A. Yost
Idaho



Dick Wallace
Vice-Chair
Washington

Tom Karier
Washington

Melinda S. Eden
Oregon

Joan M. Dukes
Oregon

December 2, 2010

MEMORANDUM

TO: Fish and Wildlife Committee

FROM: Nancy Leonard, Fish, Wildlife and Ecosystem Monitoring and Evaluation Manager

SUBJECT: ISAB's review of the draft High Level Indicator's (HLI) data and graphics and status of the draft HLI report.

ISAB Review

The ISAB provided the Council with the requested review of the data and graphics for the Council's draft High Level Indicators' Progress report (pamphlet) to Governors and Congress on November 24, 2010 (ISAB 2010-6). This review consisted of both general and specific comments for each of the three questions for which high level indicators (HLI) are used to assess progress. The ISAB review is attached to this memo (Attachment 1).

The ISAB comments provide useful suggestions for improving the HLI report. Some were incorporated in the current draft while others require more work to be addressed appropriately.

The general comments, such as including a caption for each graphic, using colors consistently, and conveying a sense of change occurring in the Basin were incorporated in the latest draft of the HLI report (Draft 11/30/10). The inclusion of target levels was done for data that have a known Council target. Trend data or historical abundance were used when available to provide context for the data. The graph captions are in development; thus, placeholders are shown on the draft. ISAB comments pertaining to the web-based version will be included when that is developed, such as providing data caveats and linking to the SOTR dam count graph.

Specific ISAB comments for each question and graph were addressed where possible in the latest draft of the HLI report (Draft 11/30/10). These comments included modifying the questions to better match the data shown, adding an ESA listed salmonid spawner abundance graph (to be included in January 2011) and a Smolt to Adult Return Ratio graph to indicate salmonid productivity.

ISAB suggestions that will require substantial work may be addressed in future editions of the HLI report. These include developing a color code system to indicate positive, negative and no change in trend data; inserting population status information on an ESU map; producing an invasive species map; adding data on the abundance of other fish species; including habitat indicators; and developing indicators that assess Program coordination.

Status of Draft HLI Report

A copy of the November 30, 2010 draft of the HLI report is provided with this packet. A PDF version is available on the Council's December 2010 Fish and Wildlife Committee agenda webpage.

Fish and Wildlife staff have worked with the Public Affairs Division to produce the text and improve the overall draft HLI report. Staff is currently working with the Public Affairs Division on the missing text for the figure captions and blank page. The text for the blank page is now in draft form and will discuss how the Fish and Wildlife Program is developed and implemented, as well as our partners' critical role in implementing the Program. In January, we will receive the compiled ESA-listed salmonid spawner abundance data from CBFWA and we will then incorporate that missing graphic.

The November 30, 2010 draft incorporates several of the ISAB suggestions to improve the HLI graphs, as discussed above.

We have shared the draft report with the entities (CBFWA, NOAA, FPC, and BPA) that have provided us with data. We are waiting to receive comments on improvements and corrections that may need to be made to the graphs.

Once we have incorporated the missing components, addressed concerns from the data providers, and made any additional changes requested by the Fish and Wildlife Committee, we will produce a revised draft for the Full Council. Once the Council is satisfied with the content and layout of the HLI report, the Public Affairs Division will finalize the draft report and produce the web-based version.

Attachment 1: ISAB Review



Independent Scientific Advisory Board
*for the Northwest Power and Conservation Council,
Columbia River Basin Indian Tribes,
and National Marine Fisheries Service*
851 SW 6th Avenue, Suite 1100
Portland, Oregon 97204

Memorandum (ISAB 2010-6)

November 24, 2010

To: Bruce Measure, Chair, Northwest Power and Conservation Council

From: Nancy Huntly, ISAB Chair

Subject: Review of the data and graphics for the Council's draft High Level Indicators' Progress Report (pamphlet) to Governors and Congress

Background

In the 2009 Columbia River Basin Fish and Wildlife Program, the Northwest Power and Conservation Council (Council) committed to adopt and periodically update high-level indicators (HLI) for the purpose of reporting program success and accomplishments to Congress, the region's governors, legislators, and citizens of the Northwest. High-level indicators will include biological, implementation, and management components. In December 2009, the Council adopted the following three questions that would be addressed in its first high level indicator report:

- 1) Are Columbia River Basin fish and wildlife abundant, diverse, productive, and spatially distributed, and sustainable?
- 2) Are mainstem hydropower dam operations meeting the Fish and Wildlife Program's survival and passage objectives?
- 3) Are Council Program actions coordinated within the Program and with other programs?

The Council's draft July 2010 Monitoring, Evaluation, Research and Reporting (MERR) Plan restates this commitment for the Council to communicate to Congress and governors on the Program's progress by using HLI. The MERR Plan further states that the Council will ask the Independent Scientific Advisory Board (ISAB) to assess the data used in reporting these HLI. This

assessment will include verifying the caveats associated with the data to ensure that any constraints are properly conveyed.

On October 1, 2010, the Council asked the ISAB to assess whether the data gathered for the 2010 HLIs are being portrayed correctly with the relevant caveats in a September 2010 draft pamphlet titled, *High Level Indicators' Data and Graphics for Reporting on Fish and Wildlife Program Progress to Governors and Congress*. The ISAB and ISRP previously reviewed the Council's proposed HLIs ([ISRP/ISAB 2009-2](#)) and the draft MERR plan ([ISAB/ISRP 2010-3](#)). This review of the draft HLI report graphics and data, and associated caveats, is a logical extension to those earlier reviews.

The ISAB understands that the draft two-page pamphlet is not in final form, and the final pamphlet will integrate a map of the Columbia River and photos of habitat, fish, and wildlife with the data-oriented graphics. The ISAB further understands that this is the Council's first attempt to assemble these data from many sources into one place. The initial effort may bring to light questions about the ease of access to data, level of effort required to achieve consistency among data sources, and remaining data gaps. The primary sources for data in the HLI report are NOAA Fisheries, the Columbia Basin Fish and Wildlife Authority (CBFWA), Fish Passage Center and Bonneville Power Administration's Pisces database. The pamphlet was organized into three sections in order to address each of the three Council questions listed above.

In addition to asking for the ISAB's assessment of whether the HLI data are being portrayed correctly with the relevant caveats, the Council also asked for the ISAB's guidance on the following points:

- Are the data used for the HLIs appropriate? Are there some HLIs for which the data are so inappropriate that the Council should not report on that HLI at this time?
- Do the caveats allow the Council to use the HLI data even though the data are not the best the Council could desire for the HLI?
- Are the graphics easy to comprehend for non-scientists?
- Do the HLIs convey the Council's message that habitat is the base of the Program?
- Input on potential titles or catchy header. Staff have suggested these thus far:
 - Habitat is the heart of the Program
 - Providing Pathways for Populations or Providing Pathways for Fish and Wildlife
 - Upstream Indicators
 - Pathways to Success

The ISAB's review below begins with general observations on the HLI graphics, and then proceeds with comments organized by the three questions posed by the Council and addressed in the two page pamphlet.

A. General observations

The ISAB recognizes the challenges of designing a brief two-page pamphlet that provides succinct, meaningful visual answers to the three high level indicator questions posed by the Council. Encapsulating the progress of such a comprehensive and complex planning and management effort as the Fish and Wildlife Program into a pamphlet is a daunting task. An effective pamphlet will require a clear focus on the questions being asked, the use of the highest quality data from monitoring programs, the use of graphics that are easily understood by the audience, selection of the best indicators for answering the questions, and a clear indication of progress to date. The graphics must be carefully designed to convey meaning with maximum clarity, compactness, consistency, and impact. As always, identifying the best data and understanding and conveying its strengths, limitations, and qualifications pose challenges. The ISAB appreciates the opportunity to provide comments on the pamphlet content and graphics at this early stage of development.

Overall, the ISAB felt that the HLI graphics as drafted only partially answered the questions posed by the Council and were often in a form that would not be easily understood by the audience. We offer some suggestions below on how to improve the clarity, consistency, and impact of the graphics.

Overall comments relevant to all figures

The use of captions – Each graphic should be designed to illustrate a specific “take home message” that the Council wants to deliver. That message should be comprehensible without too much effort by the reader. While the content and importance of the graphics used in this draft pamphlet should be obvious to many readers, an interpretive figure caption above each figure would be very helpful in clarifying the take-home message. Without this additional text, the reader may wonder what message is being delivered.

The inclusion of reference or target levels for each metric – When appropriate, each graphic should have some reference point by which to judge the data presented with reference to the question addressed. It should be *possible* to draw a conclusion, or else the graphic is not meaningful. Thus, even if the Council prefers not to lead the reader to a particular interpretation about the graphic, the graphic should show the metric relative to a specific goal, past conditions, or long-term average.

The consistent and mnemonic use of colors – The ISAB notes that although the color in figures was used to improve the attractiveness of the presentation, color was not in all cases used effectively to convey information. For example, in the draft, hatchery fish are shown in green in the salmon abundance figure and wild fish are shown in green in the harvest figure. This sort of inconsistency causes confusion and reduces readability. There should be consistent use of colors throughout the document. Secondly, there should be mnemonic use of color throughout the document, i.e., the use of colors designed to be easily remembered and associated with particular items, issues or social patterns. For example, traffic signal colors might be used to

show trends. In depicting trends as increasing (i.e., a positive effect or improving relative to Plan objectives), one might use the color green. In depicting decreasing trends or deteriorating relative to plan objectives, one might use the color red. For stable or near target conditions, one might use the color yellow. Similarly, one might use blue for wild fish (“wild blue”), metallic silver for hatchery fish (metallic = technological), and unknown hatchery/wild origin maybe a blue/metallic hatched color. Consistent use of these mnemonic colors can more effectively convey information with the great economy needed in the pamphlet.

Standardization of figure types where possible – The pamphlet should attempt to standardize or minimize the type of graphs used to present data. The existing draft uses a variety of graph types, e.g., line charts, bar charts, stacking charts that may complicate the message. For example, lamprey and sturgeon abundance data are presented in different formats.

Documentation of data sources – The online version of the pamphlet should document data sources and how the datasets were derived, plus any caveats. The pamphlet should reference the online version.

Conveying a sense of change – Although the brief pamphlet must be limited in its content, the ISAB notes that an HLI report should give the reader a sense of the rapidly changing conditions in the Columbia River Basin (e.g., climate, invasive species, and so forth). The Columbia is facing a “no-analogue” future where the processes and species shaping habitats, populations, and communities are changing rapidly and in many cases to conditions with no natural precedent. Creating and maintaining resilience and adaptive capacity in fish and wildlife populations will depend on meeting many of the goals outlined in the Fish and Wildlife Program. This important issue should be emphasized in the HLI report. Managing and preserving diverse, productive, and resilient fish and wildlife populations and their habitats will become more complex in the future. An important strategy, in addition to the habitat focus, will be to remain open to new ideas, expand cooperation, and be flexible in actions.

B. Specific comments related to questions

HLI Report Question 1: Are Columbia River Basin fish and wildlife abundant, diverse, productive, spatially distributed, and sustainable?

Four charts are used to answer this complex question. However, the four charts only address abundance of lamprey (inconsistent index counts at four dams), abundance of white sturgeon (inconsistent dataset), abundance of anadromous salmonids in 2008, and harvests of hatchery and natural origin salmonids in 2007, 2008, and 2009. Spatial distribution information presented in the graphs is confounded by missing data in some years and locations. Where the graphic is designed to reveal trends in a metric, special care is needed to ensure that the metric is comparable across years, or else the graphic may be misleading. Often, as in the case of the white sturgeon (see comments below), a subset of the data that are consistently representative will reveal trends more reliably than will an inconsistently derived aggregate of all available

data. Only indirect information is provided on productivity (through abundance) and little or no information provided on diversity, spatial structure, and sustainability. No information is provided for wildlife as a group or individual species.

Salmon figures – The 2008 graphic showing abundance of hatchery, natural origin, and unknown origin salmonids is limited because it only shows data for one year. Unfortunately, there is no time series of basinwide salmonid abundance by hatchery versus natural origin. If the goal of the chart was to state that most salmonids in the Basin are from hatchery or unknown origin, as shown in the chart, then a graphic like this would be appropriate, assuming it contained a message telling the reader how to interpret the graph. This graph would have more utility if it is linked to the CBFWA dam count figures and it is used as an example of relative abundances of hatchery, natural origin, and unknown origin salmonids. The observation that the origin of many salmonids (hatchery *versus* natural) is unknown in the harvest and spawning escapement is an important message. Inaccurate estimates of natural salmon production can complicate evaluation of stock status of these wild populations that depend on habitat whereas hatchery reared juvenile fish use the river mainly as a corridor to the estuary and ocean. Greater use of mass marking techniques in hatcheries should help to improve stock origin information. The term “natural origin salmon” is preferred over “wild salmon” in the Columbia River Basin because hatchery strays are significant in some watersheds.

The take home message in the salmon and steelhead harvest chart is not obvious. Is it important to highlight the three harvest groups (commercial, tribes, sport), or hatchery versus natural origin fish in the harvests, or is there another objective in showing these data? The chart shows harvests during three years but this short time series does not tell the reader anything about the status of natural origin or hatchery salmon, productivity, spatial distribution, sustainability, or whether harvests by these groups is stable, increasing or decreasing. Furthermore, there is no identification of species or race of salmon. This type of graph will be more important in the future when a longer time series is available. The graph is an example of how an interpretive figure caption is needed to highlight what the Council wants the viewer to see in the graph.

ISRP/ISAB ([2009-2](#)) provided a number of suggestions to address the broad question of how to convey abundance, productivity diversity, and spatial structure. They suggested that the time series of fish abundance estimates at the Columbia River mouth, Bonneville Dam, Lower Granite Dam, and Priest Rapids Dam might be used to capture both trends in abundance and spatial distribution at key locations in the Basin (see Figures by CBFWA). The overall abundance trend in the CBFWA figures are easy for readers to interpret, but they do not attempt to separate natural origin versus hatchery salmonids because the data do not exist for most years. Stacking charts such as those in the draft are not ideal because trends for each species or race of salmon are difficult to follow, but the graphs do provide a quick view of overall trends in total salmonid abundance and the viewer can gain a sense of species composition.

Another suggestion by the ISRP/ISAB to depict diversity and spatial structure was a color-coded subbasin map or ESU map (see Fig. 1 below) to indicate status trends of key natural origin

species in each subbasin or ESU, e.g., green (increasing), red (decreasing), etc. as discussed above. Such a map might focus on Chinook and/or steelhead populations since these species are the most widely distributed throughout the Basin. Some data for this effort could be obtained from the NMFS webpage (www.webapps.nwfsc.noaa.gov/apex/f?p=238:home:0), which provides some adult population data in select tributaries of select ESUs (e.g., spawner counts, sometimes fraction of spawners that are wild, sometimes productivity). ESUs that are protected by the Endangered Species Act should be indicated. However, it would take some effort to filter through this web page and develop trend indices. Alternatively, the *2008-2017 U.S. v. Oregon Management Agreement* (Pp. 28-29) identified indicator stocks that might be used to show stock-specific population trends. A relatively productive and abundant stock of fall Chinook salmon is the Hanford Reach population. Spawner estimates including hatchery strays are available for 1979-2001 for Hanford (www.critfc.org/tech/02-3report.pdf); updates should be obtainable. Ideally, these trends would focus on natural origin salmonids because natural origin salmonids are more dependent on habitat than hatchery salmonids. However, the ISAB cautions that a problem with using a small subset of stocks is that they may not be representative of the all stocks.

If this approach is not feasible, another approach to add a diversity component to the abundance graphs might be to use numbers and coded arrows by direction to show how many stocks or DPSs or ESUs are increasing, decreasing, or stable. For example, in the wild fish abundance graph, one could use something like 22 ↑ 15 ↓ 25→ to indicate that 22 stocks are increasing, 15 are decreasing and 25 stocks are stable. The values and arrows should be color coded as suggested above.




Fig. 1. As recommended by ISRP/ISAB ([2009-2](#)), a subbasin map such as this could be color-coded to indicated status trends of key species in each subbasin, e.g., green (increasing), red (decreasing), yellow (stable), white (unknown), or black (not present). The map might focus on Chinook and steelhead since these species are the most widely distributed, but the online version of the pamphlet might consider a map for each species. A map-based approach might be used to convey simple trends in other metrics, such as habitat restoration and protection. The time period covered by the trend should be noted. The influence of freshwater versus marine factors should be noted to the extent possible. See Fig. 2 also.

Productivity of salmonids also could be indicated with time series of smolt to adult survival (SARs), smolts per spawner, or adult returns per spawner. SAR data are available from Comparative Survival Study (CSS) documents. If data were available for all three metrics, then productivity in the ocean versus freshwater habitats could be identified.

Lamprey figure – The ISAB is concerned about the quality of the indices of lamprey abundance, but index counts such as these have been used and described in the status review by Close et al. (1995); they represent the best information currently available to describe lamprey status. It

may be important to point out that limitations and utility of the indices are not yet fully understood. In order to show the recent status of lamprey, which have reportedly declined over time, it would be useful to show data from earlier years, or to at least indicate the historical long-term mean index count. For example, Close et al. (1995) reported up to ~350,000 lamprey at Bonneville Dam and at the Dalles Dam in the 1960s. Missing values need to be identified on the chart.

White sturgeon figure – The chart mixes incomplete data for four different mainstem reaches, and therefore provides an inaccurate and confusing view of sturgeon abundance. If the abundance dataset for large sturgeon is used, it may be best to only use the most complete time series, i.e., abundance below Bonneville. Missing values must be indicated on the chart, otherwise viewers will misinterpret the chart. A much longer time series might be shown if harvest is used as a proxy for relative abundance, but nuances such as changing size limits must be considered. If the goal is to present information on the status of sturgeon throughout the entire Basin, then a color-coded map might be generated to highlight location-specific population trends (see comment above). Details about the trends used in this map should be described or referenced in the online version of the pamphlet.

Overall comment related to question 1

It could also be argued that because management of salmonids at this time is stock specific compared to the sturgeon, lamprey, and all other species, it may not be a good use of space to have one entire graph for sturgeon and a separate one for lamprey. Sturgeon, lamprey and *perhaps other fish and wildlife species* could be combined into one figure. For example, one could show a column of species on the left and status indicators on the right, color coded as increasing (green), decreasing (blue) or stable (yellow.). This approach (combining and perhaps expanding non-salmonids species) might allow more space for developing a map-based figure to address salmon diversity and spatial structure.

There are many species protected by the Endangered Species Act that inhabit the Columbia River Basin. The ISAB suggests a graph could show the number of ESA listed fish and wildlife species in the Basin versus the number of species that have been delisted. Colors could be used to identify species groups such as fishes, birds, mammals, amphibians, etc. This graph might also include the number of non-native invasive species that have colonized the Basin (color coded by type). The number of ESA species as well as invasive species (see ISAB 2008-4) in the Basin is quite high and this is an important message that should be delivered.

Habitat is a key component of the Fish and Wildlife Program, yet the three Council questions do not directly address habitat. Nevertheless, as noted by ISRP/ISAB (2009), the status of natural origin salmonids and wildlife is a key approach for evaluating cumulative effects of habitat quality on species of interest. It may be worthwhile to associate habitat quality to the status of species either through a statement or modification of the first question. For example, the diversity, productivity, and spatial distribution of non-hatchery salmonid populations will reflect the condition, diversity, and spatial representation of critical habitats and food webs associated

with them. If this approach is taken, then it is important to emphasize the status of natural origin salmonids rather than total salmonids, which include many hatchery fish. Natural origin salmonids depend on habitat more than hatchery salmonids. Trends in the population characteristics identified in this question could be used as a general indicator of habitat condition. However, the viewer of the pamphlet should be reminded that these population characteristics reflect habitat condition along with year to year fluctuations in the environment and harvests.

HLI Report Question 2: Are mainstem hydropower dam operations meeting the Fish and Wildlife Program's survival and passage objectives?

Question 2 is more straightforward, but the Council's survival and passage objectives need to be shown with the data so that the reader has a reference with which to compare the data. Is survival meeting the survival objectives? Historic context could be added by showing survival rates in earlier years (for spring Chinook and steelhead survival estimates in 1966-80 and 1993-99 see Williams et al. 2001 in North American Journal of Fisheries Management 21:310-317). The ISAB recommends that the graphs provide estimates of variability or confidence in the survival estimates, if possible and further notes that adult survival rates exceeding 100% will confuse readers.

The purpose of the chart showing juvenile and adult counts at each dam in 2008 is not clear. Does the Council have objectives associated with these counts, as implied by the overall question? What message is being delivered by this chart that policy makers and others will find highly informative? The juvenile count data seem to be inaccurate because they decline from 237 million fish at Lower Monumental Dam to only 7 million at McNary Dam, which includes fish from the upper Columbia.

The hydro operations are not explicitly addressed in the graphics. A more detailed analysis is needed to identify which operations produce the best results and why. Further, how have modifications to the operations impacted survival and passage? What is working well and what is not?

HLI Report Question 3: Are Council Program actions coordinated within the Program and with other programs?

Question 3 is important in that the actions of the Council's Program should be coordinated both within the Program and with other programs. However, the HLI graphics did not specifically address the question of coordinated actions – and for policy makers it will be important to do so. Instead, habitat graphics were used to address Question 3. Habitat protection and restoration is an important feature of the Fish and Wildlife Program, so charts related to habitat are important. If habitat is the focus here, then the question should be changed. Please see

ISRP/ISAB ([2009-2](#)) about caveats associated with implementation actions, such as habitat restoration, with respect to recovery of fish and wildlife species.

The graphics do not address the question about coordinated activities. However, the Fish and Wildlife staff offered the following question as an alternative: “What is the progress of implementing the Council’s Fish and Wildlife Program?” This question is much more relevant to the habitat graphics that are presented. Alternatively, this question could also be used as the overall question that is being addressed by the HLI pamphlet since trends in stock status are also a key approach for describing progress of the Program (see ISRP/ISAB [2009-2](#)). This question could be answered by 1) fish population data (existing Question 1 or something similar), 2) survival through the dams (Question 2), and 3) actions to protect and restore habitat (existing charts). For example the question might be, how much habitat has been protected and restored through implementation of the Fish and Wildlife Program? Nevertheless, if the question is altered to focus on changes in habitat conditions rather than project coordination, then the ISAB believes that it would still be worthwhile to include a graph that quantifies coordination activities over time because coordination is very important to a successful program and it implies leadership.

Context, goals, or other reference points are needed in the first two habitat graphics. For example, how much habitat has been gained relative to what is currently available? How many miles of stream have been opened up in relation to what is isolated or what is currently available? Acres of habitat purchased for fish does not seem to provide useful information by itself. Presumably these purchases secure or allow unrestricted restoration of some habitats. Stating the amount of habitat in length or area of stream and the species that will benefit from the actions might be more useful. The graphics do not make it clear if the habitat purchased information is linked to or confounded with the graphic on stream miles that are “protected.” Are the data of barriers removed shown in the second graphic linked to the graphic of stream miles made accessible? To convey the spatial aspect of the habitat actions, it would be worthwhile to show the habitat values by large region in the Basin (Fig. 2).




Fig. 2. Miles of streams protected from further hydro development in each subbasin. This map is an example of a figure type that might be used in the pamphlet to show progress with respect to habitat (although we would suggest a different color scheme, with red representing less protected habitats). The next step in this evaluation might be to calculate the percentage of “protected” stream miles in each subbasin. Map source: www.nwcouncil.org/fw/protectedareas/Default.htm

The last graphic showing wildlife habitat mitigation is much more useful than the first two graphics because the wildlife graphic provides context, i.e., wildlife habitat units lost versus the amount mitigated, and the goal for mitigation. However, most people will not be familiar with the location names shown here. A presentation of these data for several large areas of the Basin would be more informative. A graphic such as this for salmon would be useful.

C. Are the graphics easy to comprehend for non-scientists

Several comments in the section above “Overall comments relevant to all figures” directly address comprehension. Some questions may need to be modified in order to match the desired message and the data and graphics being shown. In general, the graphics in the pamphlet, as presented, would not be easy to understand by non-scientists. Fig. 3. shows a graphic that is considered to be one of the best statistical graphics of all time because it captures many quantitative values in a single, easy to interpret graphic.

D. Do the HLIs convey the Council’s message that habitat is the base of the Program?

The graphs and limited amount of text do not convey the message that habitat is the base of the Fish and Wildlife Program. ISRP/ISAB ([2009-2](#)) noted that population trends (abundance, productivity, diversity, spatial distribution) of *natural origin salmonids* (and wildlife) may be used as a metric to evaluate overall habitat availability and condition (for salmonids), but this message needs to be developed in the pamphlet before the general public will make this connection.

The habitat figures attempt to address this question, but proper context is needed for the two fish habitat graphics (see comments above). If habitat is the base of the Program, it is discouraging to see that only 20 fish passage barriers were removed in 2008, as shown in the graphic, given that there are many culverts that block fish passage in the Columbia Basin.

E. Input on potential titles or a catchy header for the report

One composite title might be:

“Pathway to recovery and sustainability”





Fig. 3. This graphic is considered by E. Tufte, an expert on visual display of quantitative information, to be one of the best statistical graphics ever drawn. The graphic shows the fate of Napoleon's army as it approached (beige band) and retreated (black band) from Moscow during 1812-1813. The graphic shows six variables, including army size across time and space, direction of army movement, and air temperature.


Columbia River Basin Fish and Wildlife Program

USING HIGH LEVEL INDICATORS TO TRACK PROGRAM PROGRESS



1 Are Columbia River Basin Fish Species Abundant, Diverse, Productive, Spatially Distributed, and Sustainable?


To understand the effectiveness of the fish and wildlife program, it is necessary to understand the status of species affected by the program. The Council addresses the status of fish populations with data for general abundance and also specific species including populations of salmon, steelhead (including the 13 ESA-listed species), Pacific lamprey, and white sturgeon. The status of the species also includes data on smolt-to-adult return ratios for salmon and steelhead, and harvest of salmon, steelhead, and white sturgeon.



Lorem ipsum dolor sit amet, consectetur adipiscing elit, sed do eiusmod tempor incididunt ut labore et dolore magna aliqua. Ut enim ad minim veniam, quis nostrud exercitation ullamco laboris nisi ut aliquip ex ea commodo consequat. Duis aute irure dolor in


ESA Salmonid Spawner Abundance

GRAPH TO BE ADDED JANUARY 2011



Smolt to Adult Return Ratio


Lorem ipsum dolor sit amet, consectetur adipiscing elit, sed do eiusmod tempor incididunt ut labore et dolore magna aliqua. Ut enim ad minim veniam, quis nostrud exercitation ullamco laboris nisi ut aliquip ex ea commodo consequat. Duis aute



Salmon and Steelhead Harvest


IN COLUMBIA RIVER MAINSTEM AND TRIBUTARIES BY FISHERY TYPE

Lorem ipsum dolor sit amet, consectetur adipiscing elit, sed do eiusmod tempor incididunt ut labore et dolore magna aliqua. Ut enim ad minim veniam, quis nostrud exercitation ullamco laboris nisi ut aliquip ex ea commodo consequat. Duis aute irure dolor in reprehenderit in voluptate velit esse cillum dolore eu fugiat nulla.




Adult Fish Abundance at Mouth of Columbia River

Lorem ipsum dolor sit amet, consectetur adipiscing elit, sed do eiusmod tempor incididunt ut labore et dolore magna aliqua. Ut enim ad minim veniam, quis nostrud exercitation ullamco laboris nisi ut aliquip ex ea commodo consequat. Duis aute irure dolor in reprehenderit in voluptate velit esse cillum dolore eu fugiat nulla.



2 Are Mainstem Hydro Operations Meeting the Council Fish and Wildlife Program's Survival and Passage Objectives?

Hydropower dams affect the survival of fish and wildlife. In fact, addressing the impacts of Columbia River Basin hydropower dams on fish and wildlife is one of the key purposes of the Northwest Power Act of 1980, the federal law that authorized the Council. To understand these impacts and the effectiveness of the program's hydropower measures, such as adult fish ladders and juvenile fish bypass systems, the Council seeks data on the survival of juvenile and adult fish passing the dams.



Lorem ipsum dolor sit amet, consectetur adipiscing elit, sed do eiusmod tempor incididunt ut labore et dolore magna aliqua. Ut enim ad minim veniam, quis nostrud exercitation ullamco laboris nisi ut aliquip ex ea commodo consequat. Duis aute irure dolor in reprehenderit in voluptate velit esse cillum dolore eu fugiat nulla.




Lorem ipsum dolor sit amet, consectetur adipiscing elit, sed do eiusmod tempor incididunt ut labore et dolore magna aliqua. Ut enim ad minim veniam, quis nostrud exercitation ullamco laboris nisi ut aliquip ex ea commodo consequat. Duis aute irure dolor in reprehenderit in voluptate velit esse cillum dolore eu fugiat nulla.


Juvenile Dam Survival Through the Columbia River Hydrosystem (Hatchery and Wild Fish)

McNary Dam to Bonneville Dam (McN-Bonn) and Lower Granite Dam to Bonneville Dam (LG-Bonn)


Lorem ipsum dolor sit amet, consectetur adipiscing elit, sed do eiusmod tempor incididunt ut labore et dolore magna aliqua. Ut enim ad minim veniam, quis nostrud exercitation ullamco



PATHWAYS FOR FISH



Fish ladders are a series of gradual steps that enable fish to swim around or over a dam. Ladders are in place at all Federal projects on the lower Columbia and lower Snake rivers.



Bypass systems are pathways made up of pipes and conduits that carry juvenile fish that enter the penstocks away from turbines and around a dam. In some cases, bypasses guide fish to collection points where they are loaded into barges or trucks and transported downstream below other dams.


3 What is the Progress of Implementing the Council's Fish and Wildlife Program?

The Council's fish and wildlife program is implemented through projects sponsored by fish and wildlife agencies, tribes, and others. To assess the effectiveness of the program it is important to understand what is being accomplished—for example, acres acquired as habitat, stream miles protected, diversion screens installed and migration barriers removed in tributaries to protect migrating juvenile fish, and habitat units (a measurement that expresses habitat quantity and quality) acquired for wildlife to compensate for losses caused by dams.

Stream Miles Made Accessible for Fish (Cumulative)

RESIDENT AND ANADROMOUS


Lorem ipsum dolor sit amet, consectetur adipiscing elit, sed do eiusmod tempor incididunt ut labore et dolore magna aliqua. Ut enim ad minim veniam, quis nostrud exercitation ullamco laboris nisi ut aliquip ex ea commodo nulla.



Barriers Removed for Fish

RESIDENT AND ANADROMOUS


Lorem ipsum dolor sit amet, consectetur adipiscing elit, sed do eiusmod tempor incididunt ut labore et dolore magna aliqua. Ut enim ad minim veniam, quis nostrud exercitation ullamco laboris nisi ut aliquip ex ea commodo nulla.



Screens Installed for Fish (Cumulative)

RESIDENT AND ANADROMOUS


Lorem ipsum dolor sit amet, consectetur adipiscing elit, sed do eiusmod tempor incididunt ut labore et dolore magna aliqua. Ut enim ad minim veniam, quis nostrud exercitation ullamco laboris nisi ut aliquip ex ea commodo nulla.



Stream Miles Protected for Wildlife and Fish Species (Cumulative)

RESIDENT AND ANADROMOUS



Lorem ipsum dolor sit amet, consectetur adipiscing elit, sed do eiusmod tempor incididunt ut labore et dolore magna aliqua. Ut enim ad minim veniam, quis nostrud exercitation ullamco laboris nisi ut aliquip ex ea.



Acres Purchased or Leased to Protect Fish Habitat (Cumulative)

RESIDENT AND ANADROMOUS

Lorem ipsum dolor sit amet, consectetur adipiscing elit, sed do eiusmod tempor incididunt ut labore et dolore magna aliqua. Ut enim ad minim veniam, quis nostrud exercitation ullamco laboris nisi ut aliquip ex ea.



Wildlife Habitat Units Lost and Mitigated per Dam

2008 STATUS


Lorem ipsum dolor sit amet, consectetur adipiscing elit, sed do eiusmod tempor incididunt ut labore et dolore magna aliqua. Ut enim ad minim veniam, quis nostrud exercitation ullamco laboris nisi ut aliquip ex ea commodo consequat. Duis aute irure dolor in reprehenderit in voluptate velit esse cillum dolore eu fugiat nulla.

Notes from the Field


Implementation of the Council's Fish and Wildlife Program relies on projects done by our Basin partners: tribal, state, federal, and local governments; private groups; and interested individuals. These partners use plans written by local groups in each tributary watershed, or "subbasin" of the Columbia River to guide selection of the most important projects to request the Bonneville Power Administration fund.




U.S. Sen. Mike Crapo, R-ID, (on the left) releasing a sockeye into Redfish Lake.



Culvert rebuilt with a natural-river bottom so that water and fish can pass between the stream and Hungry Horse Reservoir.



Wildish Track purchased to protect wildlife habitat in the Willamette River Basin, Oregon.



The Dayton Intake Dam has been modified to facilitate irrigation and reduce operating costs while protecting instream flows in the Touchet River, Dayton, Washington.

IDAHO

Snake River sockeye runs declined to the point that in 1992 a single fish returned to spawn in **Redfish Lake**. Since then, a captive broodstock program managed by the Idaho Department of Fish and Game and largely financed by the Bonneville Power Administration is preserving the species. For two decades the number of sockeye returning to Idaho fluctuated between 257 and zero, but in 2008 650 returned, last year the number was 833, and by mid-September 2010, 1,257 fish had returned.

MONTANA

The **Hungry Horse Mitigation Program**, sponsored by Montana Fish, Wildlife, and Parks, began in 1992 to address fish losses associated with the construction and operation of **Hungry Horse Dam**. The dam isolated approximately 38 percent of the Flathead Lake drainage and changed the physical and biological characteristics of the lake and river. The program's goals are to restore and reconnect critical habitat, reduce the negative interactions between native and non-native fish, and improve dam operations for native trout recovery.

OREGON

The state of Oregon and the Bonneville Power Administration signed an agreement in 2010 to jointly protect nearly 20,000 acres of wildlife habitat in the **Willamette River Basin**. Under the agreement, Bonneville purchased two conservation easements: 1,271 acres at the confluence of the Coast and Middle forks of the Willamette River, and 1,310 acres at Our Lady of Guadalupe Trappist Abbey near Lafayette. Bonneville will provide an additional \$150 million over 15 years to administer the Willamette program and to purchase an additional 16,880 acres.

WASHINGTON

Southeast Washington's Touchet River is crucial to recovery of ESA-listed steelhead. In a one-mile stretch, three water diversions were creating passage issues for steelhead, bull trout, mountain whitefish, and bridgelip suckers. Each spring instream work moved gravel and cobble berms to deliver water to the structures; and each fall these structures washed away, causing considerable damage to spawning areas. With financing from BPA, state and federal agencies, and volunteer groups, all three diversions were consolidated, ESA-compliant fish screens added, and improved fish ladders and trapping facilities built.



CONTENT
TO BE ADDED
SOON

Habitat is the path to sustainable fish and wildlife.


The Council's Fish and Wildlife Program is working to create a Columbia River Basin ecosystem that sustains abundant, productive, and diverse communities of fish and wildlife, mitigating for adverse effects caused by development and operation of the hydroelectric system. This is a long-term commitment requiring investment on the part of all Basin partners.



Fenders blue butterfly (*Icaricia icarioides fenderi*)



White sturgeon (*Acipenser transmontanus*)



McNary Dam Fish Ladder, close to the Washington side of the Columbia River




Columbia River Gorge



Collecting macro-invertebrate in Loup Creek in the Okanogan Basin.



Sockeye salmon returning to Redfish Lake, Idaho



Wildflowers along the Deschutes River, Oregon.



851 SW Sixth Avenue, Suite 1100
Portland, Oregon 97204-1348
Telephone: 503-222-5161
Toll Free: 1-800-452-5161
www.nwcouncil.org

The Northwest Power and Conservation Council works to protect, mitigate, and enhance fish and wildlife affected by the development, operation, and management of hydroelectric dams in the Columbia River Basin while assuring the Pacific Northwest an adequate, efficient, economical, and reliable power supply.

Data for this publication were provided by the Status of the Resource Report published by the Columbia River Basin Fish and Wildlife Authority, the Fish Passage Center, the National Oceanic and Atmospheric Administration, and the Bonneville Power Administration

Caveats associated with the data can be found on the Council's website:
www.nwcouncil.org
as well as on the data providers' websites:
www.sotr.cbfwa.org
www.fpc.org
www.nwr.noaa.gov/Salmon-Hydropower/Columbia-Snake-Basin/FCRPS-fish-psg.cfm
www.cbfish.org

COVER PHOTO: JEROME CHARAQUI; INSIDE MAP: LUIS PRADO; PATHWAYS FOR REHABILITATIONS: LUIS PRADO; BASED ON ILLUSTRATIONS BY BONNEVILLE POWER ADMINISTRATION; NOTES FROM THE FIELD PHOTOS FROM TOP: IDAHO: VICKY OSBORN; BIGS, MONTANA: MONTANA FISH, WILDLIFE & PARKS; OREGON: NATURE CONSERVANCY; OREGON, WASHINGTON: JADA; HABITAT PAGE PHOTOS (CLOCKWISE FROM TOP): FENDERS BUTTERFLY: GEORGE GENTRY; US FISH & WILDLIFE SERVICES; MCNARY FISH LADDER: NORTHWEST POWER AND CONSERVATION COUNCIL; GARNET FISH: GERALD COLUMBIA RIVER GORGE; JAKE BERGOT; SALMON: IDAHO FISH AND GAME; DESCHUTES RIVER: PAT FOLETT; WAT AT STEVENS: BRIAN WILLET; COASTAL TREES: PACIFIC WHITE STURGEON: OREGON DEPARTMENT OF FISH & WILDLIFE



DRAFT 11/30/10

