

Appendix List for Upper Middle Mainstem Columbia River

Appendix A. WNHP. 2003. Known High Quality or Rare Plant Communities and Wetland Ecosystems

Appendix B. IBIS. 2003. Tables of Wildlife Species Occurrence, Fish Species, T&E, Partners in Flight, and Wildlife Game Species

Appendix C. Focal Species Information: Red-winged Blackbird

Appendix D. USDA, Farm Service Agency. Conservation Reserve Program: Monthly Contract Report, Summary for Active contracts for all Program Years (1986-2004). Available: <http://www.fsa.usda.gov/crpstorpt/08Approved/rlsumyr.wa.htm>

Appendix E. Water Quality Parameters Affected by Hydropower Production

Appendix F. PNAMP. 2004. Considerations for Monitoring in Subbasin Plans from the Pacific Northwest Aquatic Monitoring Partnership.

Appendix A

Upper Middle Mainstem Columbia River Subbasin Plan

Known High Quality or Rare Plant Communities and Wetland Ecosystems

Table 1 Known high quality or rare plant communities and wetland ecosystems of the UMM Subbasin, WA.

| SCIENTIFIC NAME | COMMON NAME |
|---|--|
| <i>Abies amabilis</i> - <i>Tsuga mertensiana</i> cover type | Pacific silver fir - mountain hemlock forest |
| <i>Abies amabilis</i> / <i>Achlys triphylla</i> forest | Pacific silver fir / vanillaleaf |
| <i>Abies amabilis</i> cover type | Pacific silver fir forest |
| <i>Abies grandis</i> / <i>Acer circinatum</i> forest | Grand fir / vine maple |
| <i>Abies lasiocarpa</i> / <i>Calamagrostis rubescens</i> forest | Subalpine fir / pinegrass |
| <i>Abies lasiocarpa</i> / <i>Ledum glandulosum</i> forest | Subalpine fir / glandular labrador-tea |
| <i>Abies lasiocarpa</i> / <i>Rhododendron albiflorum</i> woodland | Subalpine fir / cascade azalea |
| <i>Abies lasiocarpa</i> / <i>Vaccinium scoparium</i> forest | Subalpine fir / grouseberry |
| <i>Abies lasiocarpa</i> cover type | Subalpine fir forest |
| <i>Abies procera</i> cover type | Noble fir forest |
| <i>Acer circinatum</i> cover type | Vine maple shrubland |
| <i>Alnus viridis</i> ssp. <i>Sinuata</i> shrubland (provisional) | Sitka alder |

| SCIENTIFIC NAME | COMMON NAME |
|--|---|
| Artemisia arbuscula / Festuca idahoensis dwarf-shrub herbaceous vegetation | Low sagebrush / Idaho fescue |
| Artemisia rigida / Poa secunda dwarf-shrub herbaceous vegetation | Stiff sagebrush / Sandberg's bluegrass |
| Artemisia rigida cover type | Stiff sagebrush shrubland |
| Artemisia tridentata / Festuca idahoensis shrub herbaceous vegetation | Big sagebrush / Idaho fescue |
| Artemisia tridentata cover type | Big sagebrush shrubland |
| Artemisia tridentata ssp. Wyomingensis / pseudoroegneria spicata shrub herbaceous vegetation | Wyoming big sagebrush / bluebunch wheatgrass |
| Artemisia tridentata ssp. Wyomingensis / stipa comata shrubland | Wyoming big sagebrush / needle-and- thread |
| Artemisia tripartita / Festuca campestris shrub herbaceous vegetation | Threetip sagebrush / rough fescue |
| Artemisia tripartita / Festuca idahoensis shrub herbaceous vegetation | Threetip sagebrush / Idaho fescue |
| Artemisia tripartita / Pseudoroegneria spicata shrub herbaceous vegetation | Threetip sagebrush / bluebunch wheatgrass |
| Artemisia tripartita / Stipa comata shrub herbaceous vegetation | Threetip sagebrush / needle-and-thread |
| Betula occidentalis / Cornus sericea shrubland | Water birch / red-osier dogwood |
| Betula occidentalis cover type | Water birch forest |
| Carex cover type | Sedge spp. Grassland |
| Carex scopulorum herbaceous vegetation | Holm's rocky mountain sedge |
| Carex utriculata herbaceous vegetation | Northwest territory sedge |
| Crataegus douglasii / Rosa woodsii | Black hawthorn / Wood's rose |

| SCIENTIFIC NAME | COMMON NAME |
|---|---|
| Danthonia intermedia herbaceous vegetation | Timber oatgrass |
| Distichlis spicata herbaceous vegetation | Saltgrass |
| Dryas octopetala dwarf-shrub herbaceous vegetation | Eight petal mountain-avens |
| Eleocharis palustris intermittently flooded herbaceous vegetation | Creeping spikerush |
| Elymus lanceolatus - Stipa comata herbaceous vegetation | Streamside wildrye - needle-and-thread |
| Eriogonum thymoides / Poa secunda dwarf-shrub herbaceous vegetation | Thyme buckwheat / Sandberg's bluegrass |
| Festuca idahoensis - Eriogonum heracleoides herbaceous vegetation | Idaho fescue - parsnip-flower buckwheat |
| Grayia spinosa / Poa secunda shrubland | Spiny hopsage / sandberg's bluegrass |
| Inland saline wetland cb | Inland saline wetland cb |
| Larix lyallii association | Subalpine larch community |
| Larix occidentalis cover type | Western larch forest |
| Leymus cinereus - Distichlis spicata herbaceous vegetation | Great basin wildrye - saltgrass |
| Philadelphus lewisii intermittently flooded shrubland | Mock orange |
| Picea engelmannii - Abies lasiocarpa cover type | Engelmann spruce - subalpine fir forest |

| SCIENTIFIC NAME | COMMON NAME |
|---|---|
| Picea engelmannii / Equisetum arvense forest | Engelmann spruce / field horsetail |
| Picea engelmannii cover type | Engelmann spruce forest |
| Pinus albicaulis - Abies lasiocarpa cover type | White-bark pine - subalpine fir forest |
| Pinus albicaulis cover type | White-bark pine forest |
| Pinus contorta cover type | Lodgepole pine forest |
| Pinus monticola cover type | Western white pine forest |
| Pinus ponderosa - Pseudotsuga menziesii / Pseudoroegneria spicata ssp. Inermis woodland | Ponderosa pine - douglas-fir / bluebunch wheatgrass |
| Pinus ponderosa - Pseudotsuga menziesii / Purshia tridentata woodland | Ponderosa pine - douglas-fir / bitterbrush |
| Pinus ponderosa - Pseudotsuga menziesii cover type | Ponderosa pine - douglas-fir forest |
| Pinus ponderosa / Calamagrostis rubescens forest | Ponderosa pine / pinegrass |
| Pinus ponderosa / Purshia tridentata woodland | Ponderosa pine / bitterbrush |
| Pinus ponderosa / Symphoricarpos albus temporarily flooded woodland | Ponderosa pine - common snowberry |
| Pinus ponderosa cover type | Ponderosa pine forest |
| Populus tremuloides / Symphoricarpos albus forest | Quaking aspen / common snowberry |
| Populus tremuloides cover type | Quaking aspen forest |

| SCIENTIFIC NAME | COMMON NAME |
|---|---|
| Pseudoroegneria spicata - Festuca Idahoensis Canyon Herbaceous Vegetation | Bluebunch wheatgrass - Idaho fescue canyon |
| Pseudoroegneria spicata - Poa secunda herbaceous vegetation | Bluebunch wheatgrass - Sandberg's bluegrass |
| Pseudoroegneria spicata cover type | Bluebunch wheatgrass grassland |
| Pseudotsuga menziesii - Abies grandis cover type | Douglas-fir - grand fir forest |
| Pseudotsuga menziesii - Abies lasiocarpa cover type | Douglas-fir - subalpine fir forest |
| Pseudotsuga menziesii - Tsuga heterophylla cover type | Douglas-fir - western hemlock forest |
| Pseudotsuga menziesii / Acer circinatum forest | Douglas-fir / vine maple |
| Pseudotsuga menziesii / Arctostaphylos uva-ursi - purshia tridentata forest | Douglas-fir / kinikinnick - bitterbrush |
| Pseudotsuga menziesii / Arctostaphylos uva-ursi cascadian forest | Douglas-fir / kinikinnick cascadian forest |
| Pseudotsuga menziesii / Calamagrostis rubescens forest | Douglas-fir / pinegrass |
| Pseudotsuga menziesii / Symphoricarpos albus forest | Douglas-fir / common snowberry |
| Pseudotsuga menziesii cover type | Douglas-fir forest |
| Purshia tridentata / Festuca idahoensis shrub herbaceous vegetation | Bitterbrush / Idaho fescue |
| Purshia tridentata / Oryzopsis hymenoides shrubland | Bitterbrush / indian ricegrass |
| Purshia tridentata / Pseudoroegneria spicata shrub herbaceous vegetation | Bitterbrush / bluebunch wheatgrass |
| Purshia tridentata / Stipa comata shrub | Bitterbrush / needle-and-thread |

| SCIENTIFIC NAME | COMMON NAME |
|---|---|
| Quercus garryana / Carex geyeri woodland | Oregon white oak / Geyer's sedge |
| Quercus garryana forest (provisional) | Oregon white oak |
| Rhus glabra / Pseudoroegneria spicata shrub herbaceous vegetation | Smooth sumac / bluebunch wheatgrass |
| Salix amygdaloides / Salix exigua woodland | Peach-leaf willow / sandbar willow |
| Salix drummondiana / Carex scopulorum var. Prionophylla shrubland | Drummond's willow / Holm's rocky mountain sedge |
| Salix planifolia / Carex scopulorum shrubland | Tea-leaf willow / Holm's rocky mountain sedge |
| Sarcobatus vermiculatus / Distichlis spicata shrubland | Greasewood / saltgrass |
| Scirpus maritimus herbaceous vegetation | Seacoast bulrush |
| Sporobolus cryptandrus - Poa secunda herbaceous vegetation | Sand dropseed - Sandberg's bluegrass |
| Stipa comata cover type | Needle-and-thread grassland |
| Subalpine freshwater wetland ec | Subalpine freshwater wetland ec |
| Subalpine riparian wetland ec | Subalpine riparian wetland ec |
| Thuja plicata - Tsuga heterophylla cover type | Western redcedar - western hemlock forest |
| Tsuga heterophylla / Mahonia nervosa var. Nervosa forest | Western hemlock / dwarf oregongrape |
| Tsuga mertensiana - Abies lasiocarpa cover type | Mountain hemlock - subalpine fir community |

| SCIENTIFIC NAME | COMMON NAME |
|------------------------|--------------------|
| Vernal pond cb | Vernal pond cb |

(WNHP 2003)

Appendix B

Upper Middle Mainstem Columbia River Subbasin

Fish and Wildlife

Table 2 Wildlife species occurrence by focal habitat type in the UMM Subbasin, WA.

| Shrubsteppe | Riparian Wetlands | Herbaceous Wetlands | Agriculture |
|---------------------------|--------------------------|----------------------------------|----------------------------------|
| American Avocet | American Badger | Tiger Salamander | Long-toed Salamander |
| American Badger | American Beaver | Northwestern Salamander | Ensatina |
| American Crow | American Crow | Long-toed Salamander | Great Basin Spadefoot |
| American Goldfinch | American Dipper | Rough-skinned Newt | Pacific Chorus (Tree) Frog |
| American Kestrel | American Goldfinch | Great Basin Spadefoot | Painted Turtle |
| American Robin | American Kestrel | Western Toad | Southern Alligator Lizard |
| Bank Swallow | | Woodhouse's Toad | Western Fence Lizard |
| Barn Owl | American Redstart | Pacific Chorus (Tree) Frog | Western Skink |
| Barn Swallow | American Robin | Cascades Frog | Rubber Boa |
| Barrow's Goldeneye | American Tree Sparrow | Columbia Spotted Frog | Racer |
| Big Brown Bat | American Wigeon | Northern Leopard Frog | Gopher Snake |
| Black Bear | Bank Swallow | Bullfrog | Western Terrestrial Garter Snake |
| Black-billed Magpie | Barn Owl | Painted Turtle | Northwestern Garter Snake |
| Black-chinned Hummingbird | Barn Swallow | Western Terrestrial Garter Snake | Common Garter Snake |
| Black-necked Stilt | Barred Owl | Common Garter Snake | Western Rattlesnake |
| Black-tailed Jackrabbit | Belted Kingfisher | Common Loon | American Bittern |
| Black-throated Sparrow | Big Brown Bat | Pied-billed Grebe | Turkey Vulture |
| Blue Grouse | Black Bear | Red-necked Grebe | Gadwall |
| Bobcat | Black Swift | Eared Grebe | American Wigeon |
| Brewer's Blackbird | Black-backed Woodpecker | Western Grebe | Mallard |
| Brewer's Sparrow | Black-billed Magpie | Clark's Grebe | Blue-winged Teal |
| Brown-headed Cowbird | Black-capped Chickadee | Double-crested Cormorant | Green-winged Teal |

| Shrubsteppe | Riparian Wetlands | Herbaceous Wetlands | Agriculture |
|---------------------------|-----------------------------|----------------------------|----------------------|
| Bullfrog | Black-chinned Hummingbird | American Bittern | Northern Harrier |
| Burrowing Owl | Black-crowned Night-heron | Great Blue Heron | Swainson's Hawk |
| Bushy-tailed Woodrat | Black-headed Grosbeak | Great Egret | Red-tailed Hawk |
| California Myotis | Black-tailed Deer | Black-crowned Night-heron | Ferruginous Hawk |
| California Quail | Black-throated Gray Warbler | Turkey Vulture | American Kestrel |
| Canada Goose | Blue Grouse | Canada Goose | Prairie Falcon |
| Canyon Wren | Bobcat | Tundra Swan | Chukar |
| Chipping Sparrow | Bobolink | Wood Duck | Gray Partridge |
| Chukar | Bohemian Waxwing | Gadwall | Ring-necked Pheasant |
| Cliff Swallow | Brewer's Blackbird | American Wigeon | Ruffed Grouse |
| Columbia Spotted Frog | Brown Creeper | Mallard | Sage Grouse |
| Columbian Ground Squirrel | Brown-headed Cowbird | Blue-winged Teal | Sharp-tailed Grouse |
| Common Garter Snake | Bullfrog | Cinnamon Teal | Wild Turkey |
| Common Nighthawk | Bullock's Oriole | Northern Shoveler | California Quail |
| Common Poorwill | Bushy-tailed Woodrat | Northern Pintail | Virginia Rail |
| Common Porcupine | California Myotis | Green-winged Teal | Sora |
| Common Raven | California Quail | Canvasback | American Coot |
| Cooper's Hawk | Calliope Hummingbird | Redhead | Killdeer |
| Coyote | Canada Goose | Ring-necked Duck | Black-necked Stilt |
| Deer Mouse | Canyon Wren | Barrow's Goldeneye | American Avocet |
| Eastern Kingbird | Cascade Frog | Hooded Merganser | Long-billed Curlew |
| European Starling | Cassin's Finch | Ruddy Duck | Wilson's Snipe |
| Ferruginous Hawk | Cassin's Vireo | Northern Harrier | Ring-billed Gull |
| Fringed Myotis | Cedar Waxwing | Sharp-shinned Hawk | Rock Dove |
| Golden Eagle | Chipping Sparrow | Cooper's Hawk | Mourning Dove |
| Gopher Snake | Chukar | Northern Goshawk | Barn Owl |
| Grasshopper Sparrow | Cliff Swallow | Swainson's Hawk | Western Screech-owl |
| Gray Flycatcher | Coast Mole | Red-tailed Hawk | Great Horned Owl |
| Gray Partridge | Columbia Spotted Frog | Rough-legged Hawk | Burrowing Owl |
| Great Basin Pocket Mouse | Columbian Ground Squirrel | Golden Eagle | Long-eared Owl |

| Shrubsteppe | Riparian Wetlands | Herbaceous Wetlands | Agriculture |
|-------------------------------|--------------------------|----------------------------|---------------------------|
| Great Basin Spadefoot | Columbian Mouse | American Kestrel | Short-eared Owl |
| Great Horned Owl | Common Garter Snake | Gyrfalcon | Common Nighthawk |
| Greater Yellowlegs | Common Merganser | Ring-necked Pheasant | Common Poorwill |
| Hoary Bat | Common Nighthawk | Virginia Rail | Black-chinned Hummingbird |
| Horned Lark | Common Porcupine | Sora | Rufous Hummingbird |
| Killdeer | Common Raven | American Coot | Lewis's Woodpecker |
| Lark Sparrow | Common Redpoll | Killdeer | Red-breasted Sapsucker |
| Least Chipmunk | Common Yellowthroat | Black-necked Stilt | Downy Woodpecker |
| Lesser Yellowlegs | Cooper's Hawk | American Avocet | Hairy Woodpecker |
| Little Brown Myotis | Cordilleran Flycatcher | Greater Yellowlegs | Northern Flicker |
| Loggerhead Shrike | Coyote | Lesser Yellowlegs | Western Wood-pewee |
| Long-billed Curlew | Creeping Vole | Solitary Sandpiper | Willow Flycatcher |
| Long-eared Myotis | Dark-eyed Junco | Spotted Sandpiper | Say's Phoebe |
| Long-eared Owl | Deer Mouse | Long-billed Curlew | Western Kingbird |
| Long-legged Myotis | Double-crested Cormorant | Western Sandpiper | Eastern Kingbird |
| Long-tailed Vole | Downy Woodpecker | Least Sandpiper | Loggerhead Shrike |
| Long-tailed Weasel | Dusky Flycatcher | Baird's Sandpiper | Warbling Vireo |
| Long-toed Salamander | Eastern Cottontail | Pectoral Sandpiper | Steller's Jay |
| Mallard | Eastern Fox Squirrel | Long-billed Dowitcher | Black-billed Magpie |
| Merriam's Shrew | Eastern Kingbird | Wilson's Snipe | American Crow |
| Mink | Ermine | Wilson's Phalarope | Common Raven |
| Montane Vole | European Starling | Ring-billed Gull | Horned Lark |
| Mountain Bluebird | Evening Grosbeak | California Gull | Tree Swallow |
| Mourning Dove | | Herring Gull | Violet-green Swallow |
| Nashville Warbler | | Thayer's Gull | Cliff Swallow |
| Night Snake | Fox Sparrow | Glaucous Gull | Barn Swallow |
| Northern Flicker | Fringed Myotis | Caspian Tern | Black-capped Chickadee |
| Northern Goshawk | Golden Eagle | Forster's Tern | White-breasted Nuthatch |
| Northern Harrier | Golden-crowned Kinglet | Black Tern | Brown Creeper |
| Northern Pocket Gopher | Gopher Snake | Western Screech-owl | House Wren |
| Northern Rough-winged Swallow | Gray Catbird | Great Horned Owl | Western Bluebird |

| Shrubsteppe | Riparian Wetlands | Herbaceous Wetlands | Agriculture |
|---------------------------------|--------------------------|-------------------------------|-----------------------------|
| Northern Shrike | Gray Jay | Snowy Owl | Mountain Bluebird |
| Nuttall's (Mountain) Cottontail | Great Basin Spadefoot | Northern Pygmy-owl | Swainson's Thrush |
| Orange-crowned Warbler | Great Blue Heron | Burrowing Owl | American Robin |
| Osprey | Great Egret | Great Gray Owl | Gray Catbird |
| Pacific Chorus (Tree) Frog | Great Horned Owl | Long-eared Owl | European Starling |
| Painted Turtle | Greater Yellowlegs | Short-eared Owl | Cedar Waxwing |
| Pallid Bat | Green-winged Teal | Common Nighthawk | Orange-crowned Warbler |
| Prairie Falcon | Hairy Woodpecker | Black Swift | Nashville Warbler |
| Pygmy Rabbit | Heather Vole | Vaux's Swift | Black-throated Gray Warbler |
| Racer | Hermit Thrush | White-throated Swift | Macgillivray's Warbler |
| Red-tailed Hawk | Hoary Bat | Black-chinned Hummingbird | Common Yellowthroat |
| Ringneck Snake | Hooded Merganser | Calliope Hummingbird | Wilson's Warbler |
| Ring-necked Pheasant | House Finch | Rufous Hummingbird | Yellow-breasted Chat |
| Rock Dove | House Wren | Eastern Kingbird | Spotted Towhee |
| Rock Wren | Killdeer | Loggerhead Shrike | Chipping Sparrow |
| Rough-legged Hawk | Lazuli Bunting | Northern Shrike | Brewer's Sparrow |
| Rough-skinned Newt | Least Chipmunk | Black-billed Magpie | Vesper Sparrow |
| Rubber Boa | Lesser Yellowlegs | American Crow | Savannah Sparrow |
| Sage Grouse | Lewis's Woodpecker | Common Raven | Grasshopper Sparrow |
| Sage Sparrow | Lincoln's Sparrow | Tree Swallow | Song Sparrow |
| Sage Thrasher | Little Brown Myotis | Violet-green Swallow | White-crowned Sparrow |
| Sagebrush Lizard | Long-eared Myotis | Northern Rough-winged Swallow | Dark-eyed Junco |
| Sagebrush Vole | Long-eared Owl | Bank Swallow | Black-headed Grosbeak |
| Savannah Sparrow | Long-legged Myotis | Cliff Swallow | Lazuli Bunting |
| Say's Phoebe | Long-tailed Vole | Barn Swallow | Bobolink |
| Sharp-shinned Hawk | Long-tailed Weasel | Black-capped Chickadee | Red-winged Blackbird |
| Sharp-tailed Grouse | Long-toed Salamander | Marsh Wren | Western Meadowlark |
| Short-eared Owl | Macgillivray's Warbler | American Dipper | Yellow-headed Blackbird |
| Short-horned Lizard | Mallard | Ruby-crowned Kinglet | Brewer's Blackbird |

| Shrubsteppe | Riparian Wetlands | Herbaceous Wetlands | Agriculture |
|----------------------------------|-------------------------------|-----------------------------|---------------------------------|
| Side-blotched Lizard | Masked Shrew | American Robin | Brown-headed Cowbird |
| Snow Bunting | Meadow Vole | European Starling | Bullock's Oriole |
| Solitary Sandpiper | Mink | American Pipit | House Finch |
| Spotted Bat | Montane Shrew | Cedar Waxwing | American Goldfinch |
| Spotted Sandpiper | Montane Vole | Yellow-rumped Warbler | House Sparrow |
| Striped Whipsnake | Moose | Common Yellowthroat | Vagrant Shrew |
| Swainson's Hawk | Mountain Bluebird | Savannah Sparrow | Trowbridge's Shrew |
| Tiger Salamander | Mountain Chickadee | Song Sparrow | Shrew-mole |
| Townsend's Big-eared Bat | Mountain Lion | Lincoln's Sparrow | Coast Mole |
| Townsend's Ground Squirrel | Mourning Dove | White-crowned Sparrow | California Myotis |
| Townsend's Solitaire | Muskrat | Lapland Longspur | Yuma Myotis |
| Turkey Vulture | Nashville Warbler | Bobolink | Little Brown Myotis |
| Vagrant Shrew | Northern Alligator Lizard | Red-winged Blackbird | Long-legged Myotis |
| Vesper Sparrow | Northern Flicker | Western Meadowlark | Fringed Myotis |
| Washington Ground Squirrel | Northern Flying Squirrel | Yellow-headed Blackbird | Long-eared Myotis |
| Western Fence Lizard | Northern Goshawk | Brewer's Blackbird | Big Brown Bat |
| Western Harvest Mouse | Northern Harrier | Brown-headed Cowbird | Spotted Bat |
| Western Kingbird | | House Finch | Townsend's Big-eared Bat |
| Western Meadowlark | Northern Pocket Gopher | Pine Siskin | Pallid Bat |
| Western Pipistrelle | Northern Pygmy-owl | American Goldfinch | Eastern Cottontail |
| Western Rattlesnake | Northern River Otter | Vagrant Shrew | Nuttall's (Mountain) Cottontail |
| Western Skink | Northern Rough-winged Swallow | Pacific Water Shrew | Snowshoe Hare |
| Western Small-footed Myotis | Northern Saw-whet Owl | Shrew-mole | White-tailed Jackrabbit |
| Western Terrestrial Garter Snake | Northern Waterthrush | California Myotis | Black-tailed Jackrabbit |
| Western Toad | Northwestern Salamander | Western Small-footed Myotis | Least Chipmunk |
| White-crowned Sparrow | Olive-sided Flycatcher | Yuma Myotis | Yellow-bellied Marmot |
| White-tailed Jackrabbit | Orange-crowned Warbler | Little Brown Myotis | Washington Ground Squirrel |

| Shrubsteppe | Riparian Wetlands | Herbaceous Wetlands | Agriculture |
|-----------------------|----------------------------|----------------------------|----------------------------|
| White-throated Swift | Osprey | Long-legged Myotis | Columbian Ground Squirrel |
| Woodhouse's Toad | Pacific Chorus (Tree) Frog | Fringed Myotis | Eastern Fox Squirrel |
| Yellow-bellied Marmot | Pacific Jumping Mouse | Long-eared Myotis | Northern Pocket Gopher |
| Yuma Myotis | Pacific Water Shrew | Silver-haired Bat | Great Basin Pocket Mouse |
| Mule deer | Painted Turtle | Big Brown Bat | Western Harvest Mouse |
| Elk | Pallid Bat | Hoary Bat | Deer Mouse |
| | Pied-billed Grebe | Spotted Bat | Northern Grasshopper Mouse |
| | Pileated Woodpecker | Townsend's Big-eared Bat | Bushy-tailed Woodrat |
| | Pine Siskin | Pallid Bat | Montane Vole |
| | Prairie Falcon | Yellow-bellied Marmot | Long-tailed Vole |
| | Pygmy Nuthatch | American Beaver | Creeping Vole |
| | Raccoon | Western Harvest Mouse | Muskrat |
| | Racer | Deer Mouse | Black Rat |
| | Red Crossbill | Meadow Vole | Norway Rat |
| | Red Fox | Montane Vole | House Mouse |
| | Red-breasted Nuthatch | Long-tailed Vole | Western Jumping Mouse |
| | Red-breasted Sapsucker | Muskrat | Pacific Jumping Mouse |
| | Red-eyed Vireo | Northern Bog Lemming | Coyote |
| | Red-naped Sapsucker | Western Jumping Mouse | Red Fox |
| | Red-tailed Hawk | Pacific Jumping Mouse | Raccoon |
| | Red-winged Blackbird | Common Porcupine | Ermine |
| | Ring-necked Duck | Nutria | Long-tailed Weasel |
| | Ring-necked Pheasant | Coyote | American Badger |
| | Rough-legged Hawk | Black Bear | Striped Skunk |
| | Rough-skinned Newt | Grizzly Bear | Bobcat |
| | Rubber Boa | Raccoon | Rocky Mountain Elk |
| | Ruby-crowned Kinglet | Long-tailed Weasel | |
| | Ruffed Grouse | Mink | |
| | Rufous Hummingbird | Striped Skunk | |
| | Savannah Sparrow | Northern River Otter | |
| | Say's Phoebe | Mountain Lion | |

| Shrubsteppe | Riparian Wetlands | Herbaceous Wetlands | Agriculture |
|--------------------|---------------------------|----------------------------|--------------------|
| | Sharp-tailed Grouse | Bobcat | |
| | Shrew-mole | Rocky Mountain Elk | |
| | Silver-haired Bat | Mule Deer | |
| | Snowshoe Hare | White-tailed Deer | |
| | Solitary Sandpiper | | |
| | Song Sparrow | | |
| | Southern Alligator Lizard | | |
| | Southern Red-backed Vole | | |
| | Spotted Bat | | |
| | Spotted Sandpiper | | |
| | Spotted Towhee | | |
| | Steller's Jay | | |
| | Striped Skunk | | |
| | Swainson's Hawk | | |
| | Swainson's Thrush | | |
| | Tailed Frog | | |
| | Three-toed Woodpecker | | |
| | Tiger Salamander | | |
| | Townsend's Big-eared Bat | | |
| | Townsend's Solitaire | | |
| | Townsend's Warbler | | |
| | Tree Swallow | | |
| | Trowbridge's Shrew | | |
| | Turkey Vulture | | |
| | Vagrant Shrew | | |
| | Vaux's Swift | | |
| | Veery | | |
| | Violet-green Swallow | | |
| | Warbling Vireo | | |
| | Water Shrew | | |
| | Water Vole | | |
| | Western Bluebird | | |

| Shrubsteppe | Riparian Wetlands | Herbaceous Wetlands | Agriculture |
|--------------------|----------------------------------|----------------------------|--------------------|
| | Western Harvest Mouse | | |
| | Western Jumping Mouse | | |
| | Western Pipistrelle | | |
| | Western Rattlesnake | | |
| | Western Screech-owl | | |
| | Western Small-footed Myotis | | |
| | Western Tanager | | |
| | Western Terrestrial Garter Snake | | |
| | Western Toad | | |
| | Western Wood-pewee | | |
| | White-breasted Nuthatch | | |
| | White-crowned Sparrow | | |
| | White-headed Woodpecker | | |
| | White-tailed Jackrabbit | | |
| | White-throated Swift | | |
| | Wild Turkey | | |
| | Williamson's Sapsucker | | |
| | Willow Flycatcher | | |
| | Wilson's Warbler | | |
| | Winter Wren | | |
| | Wood Duck | | |
| | Woodhouse's Toad | | |
| | Yellow Warbler | | |
| | Yellow-bellied Marmot | | |
| | Yellow-breasted Chat | | |
| | Yellow-pine Chipmunk | | |
| | Yellow-rumped Warbler | | |
| | Yuma Myotis | | |

(IBIS 2003)

Table 3 Wildlife species occurrence for the UMM Subbasin

| | Common Name | Scientific Name | Salmonid Relationship | Closely Associated with Riparian Wetlands | Closely Associated with Other Wetlands |
|-------------------|-------------------------------|--------------------------|------------------------------|--|---|
| Amphibians | | | | | |
| | Tiger Salamander | Ambystoma tigrinum | | 1 | |
| | Northwestern Salamander | Ambystoma gracile | | | |
| | Long-toed Salamander | Ambystoma macrodactylum | | 1 | |
| | Pacific Giant Salamander | Dicamptodon tenebrosus | 1 | | |
| | Rough-skinned Newt | Taricha granulosa | | | 1 |
| | Western Red-backed Salamander | Plethodon vehiculum | | | |
| | Ensatina | Ensatina eschscholtzii | | | |
| | Tailed Frog | Ascaphus truei | | 1 | |
| | Great Basin Spadefoot | Scaphiopus intermontanus | | 1 | |
| | Western Toad | Bufo boreas | | 1 | |
| | Woodhouse's Toad | Bufo woodhousii | | 1 | |
| | Pacific Chorus (Tree) Frog | Pseudacris regilla | | 1 | |
| | Cascades Frog | Rana cascadae | | | |
| | Columbia Spotted Frog | Rana luteiventris | | 1 | |
| | Bullfrog | Rana catesbeiana | | 1 | |
| | Total Amphibians: | 15 | Total: 1 | 9 | 1 |
| Birds | | | | | |
| | Common Loon | Gavia immer | 1 | | 1 |
| | Pied-billed Grebe | Podilymbus podiceps | 1 | | 1 |
| | Red-necked Grebe | Podiceps grisegena | 1 | | 1 |
| | Eared Grebe | Podiceps nigricollis | | | 1 |

| | Common Name | Scientific Name | Salmonid Relationship | Closely Associated with Riparian Wetlands | Closely Associated with Other Wetlands |
|--|---------------------------|----------------------------------|------------------------------|--|---|
| | Western Grebe | <i>Aechmophorus occidentalis</i> | 1 | | 1 |
| | Clark's Grebe | <i>Aechmophorus clarkii</i> | 1 | | 1 |
| | Double-crested Cormorant | <i>Phalacrocorax auritus</i> | 1 | 1 | |
| | American Bittern | <i>Botaurus lentiginosus</i> | | | 1 |
| | Great Blue Heron | <i>Ardea herodias</i> | 1 | 1 | |
| | Great Egret | <i>Ardea alba</i> | 1 | 1 | |
| | Black-crowned Night-heron | <i>Nycticorax nycticorax</i> | 1 | 1 | |
| | Turkey Vulture | <i>Cathartes aura</i> | 1 | | |
| | Canada Goose | <i>Branta canadensis</i> | | | 1 |
| | Tundra Swan | <i>Cygnus columbianus</i> | | | |
| | Wood Duck | <i>Aix sponsa</i> | | 1 | |
| | Gadwall | <i>Anas strepera</i> | | | 1 |
| | American Wigeon | <i>Anas americana</i> | | | 1 |
| | Mallard | <i>Anas platyrhynchos</i> | 1 | 1 | |
| | Blue-winged Teal | <i>Anas discors</i> | | | 1 |
| | Cinnamon Teal | <i>Anas cyanoptera</i> | | | 1 |
| | Northern Shoveler | <i>Anas clypeata</i> | | | 1 |
| | Northern Pintail | <i>Anas acuta</i> | | | 1 |
| | Green-winged Teal | <i>Anas crecca</i> | 1 | | 1 |
| | Canvasback | <i>Aythya valisineria</i> | 1 | | 1 |
| | Redhead | <i>Aythya americana</i> | | | 1 |
| | Ring-necked Duck | <i>Aythya collaris</i> | | | |
| | Greater Scaup | <i>Aythya marila</i> | 1 | | |
| | Barrow's Goldeneye | <i>Bucephala islandica</i> | 1 | | |
| | Hooded Merganser | <i>Lophodytes cucullatus</i> | 1 | 1 | |
| | Common Merganser | <i>Mergus merganser</i> | 1 | 1 | |

| | Common Name | Scientific Name | Salmonid Relationship | Closely Associated with Riparian Wetlands | Closely Associated with Other Wetlands |
|--|----------------------|----------------------------------|------------------------------|--|---|
| | Ruddy Duck | <i>Oxyura jamaicensis</i> | | | 1 |
| | Osprey | <i>Pandion haliaetus</i> | 1 | | |
| | Northern Harrier | <i>Circus cyaneus</i> | | | |
| | Sharp-shinned Hawk | <i>Accipiter striatus</i> | | | |
| | Cooper's Hawk | <i>Accipiter cooperii</i> | | | |
| | Northern Goshawk | <i>Accipiter gentilis</i> | | | |
| | Swainson's Hawk | <i>Buteo swainsoni</i> | | | |
| | Red-tailed Hawk | <i>Buteo jamaicensis</i> | 1 | | |
| | Ferruginous Hawk | <i>Buteo regalis</i> | | | |
| | Rough-legged Hawk | <i>Buteo lagopus</i> | | | |
| | Golden Eagle | <i>Aquila chrysaetos</i> | 1 | | |
| | American Kestrel | <i>Falco sparverius</i> | | | |
| | Gyr Falcon | <i>Falco rusticolus</i> | 1 | | |
| | Prairie Falcon | <i>Falco mexicanus</i> | | | |
| | Chukar | <i>Alectoris chukar</i> | | | |
| | Gray Partridge | <i>Perdix perdix</i> | | | |
| | Ring-necked Pheasant | <i>Phasianus colchicus</i> | | 1 | |
| | Ruffed Grouse | <i>Bonasa umbellus</i> | | 1 | |
| | Sage Grouse | <i>Centrocercus urophasianus</i> | | | |
| | Spruce Grouse | <i>Falcapennis canadensis</i> | | | |
| | Blue Grouse | <i>Dendragapus obscurus</i> | | 1 | |
| | Sharp-tailed Grouse | <i>Tympanuchus phasianellus</i> | | 1 | |
| | Wild Turkey | <i>Meleagris gallopavo</i> | | | |
| | California Quail | <i>Callipepla californica</i> | | | |
| | Virginia Rail | <i>Rallus limicola</i> | | | 1 |
| | Sora | <i>Porzana carolina</i> | | | 1 |
| | American Coot | <i>Fulica americana</i> | | | 1 |

| | Common Name | Scientific Name | Salmonid Relationship | Closely Associated with Riparian Wetlands | Closely Associated with Other Wetlands |
|--|------------------------|-------------------------|------------------------------|--|---|
| | Killdeer | Charadrius vociferus | 1 | | |
| | Black-necked Stilt | Himantopus mexicanus | | | 1 |
| | American Avocet | Recurvirostra americana | | | 1 |
| | Greater Yellowlegs | Tringa melanoleuca | 1 | | |
| | Lesser Yellowlegs | Tringa flavipes | | | |
| | Solitary Sandpiper | Tringa solitaria | | 1 | |
| | Spotted Sandpiper | Actitis macularia | 1 | | |
| | Long-billed Curlew | Numenius americanus | | | |
| | Semipalmated Sandpiper | Calidris pusilla | | | |
| | Western Sandpiper | Calidris mauri | | | |
| | Least Sandpiper | Calidris minutilla | | | |
| | Baird's Sandpiper | Calidris bairdii | | | |
| | Pectoral Sandpiper | Calidris melanotos | | | |
| | Stilt Sandpiper | Calidris himantopus | | | |
| | Long-billed Dowitcher | Limnodromus scolopaceus | | | |
| | Common Snipe | Gallinago gallinago | | | 1 |
| | Wilson's Phalarope | Phalaropus tricolor | | | 1 |
| | Red-necked Phalarope | Phalaropus lobatus | | | |
| | Ring-billed Gull | Larus delawarensis | 1 | | |
| | California Gull | Larus californicus | 1 | | |
| | Herring Gull | Larus argentatus | 1 | | |
| | Thayer's Gull | Larus thayeri | 1 | | |
| | Glaucous Gull | Larus hyperboreus | 1 | | |
| | Caspian Tern | Sterna caspia | 1 | | |
| | Forster's Tern | Sterna forsteri | 1 | | 1 |
| | Black Tern | Chlidonias niger | | | 1 |
| | Rock Dove | Columba livia | | | |

| | Common Name | Scientific Name | Salmonid Relationship | Closely Associated with Riparian Wetlands | Closely Associated with Other Wetlands |
|--|---------------------------|--------------------------|------------------------------|--|---|
| | Mourning Dove | Zenaida macroura | | 1 | |
| | Barn Owl | Tyto alba | | | |
| | Western Screech-owl | Otus kennicottii | | 1 | |
| | Great Horned Owl | Bubo virginianus | | | |
| | Snowy Owl | Nyctea scandiaca | 1 | | |
| | Northern Pygmy-owl | Glaucidium gnoma | | | |
| | Burrowing Owl | Athene cunicularia | | | |
| | Barred Owl | Strix varia | | | |
| | Great Gray Owl | Strix nebulosa | | | |
| | Long-eared Owl | Asio otus | | 1 | |
| | Short-eared Owl | Asio flammeus | | | 1 |
| | Boreal Owl | Aegolius funereus | | | |
| | Northern Saw-whet Owl | Aegolius acadicus | | | |
| | Common Nighthawk | Chordeiles minor | | | |
| | Common Poorwill | Phalaenoptilus nuttallii | | | |
| | Black Swift | Cypseloides niger | | | |
| | Vaux's Swift | Chaetura vauxi | | | |
| | White-throated Swift | Aeronautes saxatalis | | | |
| | Black-chinned Hummingbird | Archilochus alexandri | | | |
| | Calliope Hummingbird | Stellula calliope | | | |
| | Rufous Hummingbird | Selasphorus rufus | | | |
| | Belted Kingfisher | Ceryle alcyon | 1 | 1 | |
| | Lewis's Woodpecker | Melanerpes lewis | | | |
| | Williamson's Sapsucker | Sphyrapicus thyroideus | | | |

| | Common Name | Scientific Name | Salmonid Relationship | Closely Associated with Riparian Wetlands | Closely Associated with Other Wetlands |
|--|--------------------------|------------------------|------------------------------|--|---|
| | Red-naped Sapsucker | Sphyrapicus nuchalis | | 1 | |
| | Red-breasted Sapsucker | Sphyrapicus ruber | | | |
| | Downy Woodpecker | Picoides pubescens | | | |
| | Hairy Woodpecker | Picoides villosus | | | |
| | White-headed Woodpecker | Picoides albolarvatus | | | |
| | Three-toed Woodpecker | Picoides tridactylus | | | |
| | Black-backed Woodpecker | Picoides arcticus | | | |
| | Northern Flicker | Colaptes auratus | | | |
| | Pileated Woodpecker | Dryocopus pileatus | | | |
| | Olive-sided Flycatcher | Contopus cooperi | | | |
| | Western Wood-pewee | Contopus sordidulus | | | |
| | Willow Flycatcher | Empidonax traillii | 1 | 1 | |
| | Hammond's Flycatcher | Empidonax hammondii | | | |
| | Gray Flycatcher | Empidonax wrightii | | | |
| | Dusky Flycatcher | Empidonax oberholseri | | | |
| | Pacific-slope Flycatcher | Empidonax difficilis | | | |
| | Cordilleran Flycatcher | Empidonax occidentalis | | 1 | |
| | Say's Phoebe | Sayornis saya | | | |
| | Western Kingbird | Tyrannus verticalis | | | |
| | Eastern Kingbird | Tyrannus tyrannus | | | |
| | Loggerhead Shrike | Lanius ludovicianus | | | |
| | Northern Shrike | Lanius excubitor | | | |
| | Cassin's Vireo | Vireo cassinii | | | |

| | Common Name | Scientific Name | Salmonid Relationship | Closely Associated with Riparian Wetlands | Closely Associated with Other Wetlands |
|--|-------------------------------|----------------------------|------------------------------|--|---|
| | Warbling Vireo | Vireo gilvus | | 1 | |
| | Red-eyed Vireo | Vireo olivaceus | | 1 | |
| | Gray Jay | Perisoreus canadensis | 1 | | |
| | Steller's Jay | Cyanocitta stelleri | 1 | | |
| | Clark's Nutcracker | Nucifraga columbiana | | | |
| | Black-billed Magpie | Pica pica | 1 | 1 | |
| | American Crow | Corvus brachyrhynchos | 1 | | |
| | Northwestern Crow | Corvus caurinus | 1 | | |
| | Common Raven | Corvus corax | 1 | | |
| | Horned Lark | Eremophila alpestris | | | |
| | Tree Swallow | Tachycineta bicolor | 1 | 1 | |
| | Violet-green Swallow | Tachycineta thalassina | 1 | | |
| | Northern Rough-winged Swallow | Stelgidopteryx serripennis | 1 | 1 | |
| | Bank Swallow | Riparia riparia | 1 | 1 | |
| | Cliff Swallow | Petrochelidon pyrrhonota | 1 | 1 | |
| | Barn Swallow | Hirundo rustica | 1 | 1 | |
| | Black-capped Chickadee | Poecile atricapillus | | | |
| | Mountain Chickadee | Poecile gambeli | | | |
| | Chestnut-backed Chickadee | Poecile rufescens | | | |
| | Boreal Chickadee | Poecile hudsonicus | | | |
| | Red-breasted Nuthatch | Sitta canadensis | | | |
| | White-breasted Nuthatch | Sitta carolinensis | | | |
| | Pygmy Nuthatch | Sitta pygmaea | | 1 | |
| | Brown Creeper | Certhia americana | | | |

| | Common Name | Scientific Name | Salmonid Relationship | Closely Associated with Riparian Wetlands | Closely Associated with Other Wetlands |
|--|------------------------|-------------------------|------------------------------|--|---|
| | Rock Wren | Salpinctes obsoletus | | | |
| | Canyon Wren | Catherpes mexicanus | | | |
| | House Wren | Troglodytes aedon | | | |
| | Winter Wren | Troglodytes troglodytes | 1 | | |
| | Marsh Wren | Cistothorus palustris | | | 1 |
| | American Dipper | Cinclus mexicanus | 1 | 1 | |
| | Golden-crowned Kinglet | Regulus satrapa | | | |
| | Ruby-crowned Kinglet | Regulus calendula | | | |
| | Western Bluebird | Sialia mexicana | | | |
| | Mountain Bluebird | Sialia currucoides | | | |
| | Townsend's Solitaire | Myadestes townsendi | | | |
| | Veery | Catharus fuscescens | | 1 | |
| | Swainson's Thrush | Catharus ustulatus | | | |
| | Hermit Thrush | Catharus guttatus | | | |
| | American Robin | Turdus migratorius | 1 | | |
| | Varied Thrush | Ixoreus naevius | 1 | | |
| | Gray Catbird | Dumetella carolinensis | | 1 | |
| | Northern Mockingbird | Mimus polyglottos | | | |
| | Sage Thrasher | Oreoscoptes montanus | | | |
| | European Starling | Sturnus vulgaris | | 1 | |
| | American Pipit | Anthus rubescens | | | |
| | Bohemian Waxwing | Bombycilla garrulus | | | |
| | Cedar Waxwing | Bombycilla cedrorum | | 1 | |
| | Orange-crowned Warbler | Vermivora celata | | | |
| | Nashville Warbler | Vermivora ruficapilla | | | |

| | Common Name | Scientific Name | Salmonid Relationship | Closely Associated with Riparian Wetlands | Closely Associated with Other Wetlands |
|--|-----------------------------|----------------------------------|------------------------------|--|---|
| | Yellow Warbler | <i>Dendroica petechia</i> | | 1 | |
| | Yellow-rumped Warbler | <i>Dendroica coronata</i> | | | |
| | Black-throated Gray Warbler | <i>Dendroica nigrescens</i> | | | |
| | Townsend's Warbler | <i>Dendroica townsendi</i> | | | |
| | Hermit Warbler | <i>Dendroica occidentalis</i> | | | |
| | American Redstart | <i>Setophaga ruticilla</i> | | 1 | |
| | Northern Waterthrush | <i>Seiurus noveboracensis</i> | | 1 | |
| | Macgillivray's Warbler | <i>Oporornis tolmiei</i> | | | |
| | Common Yellowthroat | <i>Geothlypis trichas</i> | | 1 | |
| | Wilson's Warbler | <i>Wilsonia pusilla</i> | | | |
| | Yellow-breasted Chat | <i>Icteria virens</i> | | 1 | |
| | Western Tanager | <i>Piranga ludoviciana</i> | | | |
| | Spotted Towhee | <i>Pipilo maculatus</i> | 1 | | |
| | American Tree Sparrow | <i>Spizella arborea</i> | | | |
| | Chipping Sparrow | <i>Spizella passerina</i> | | | |
| | Brewer's Sparrow | <i>Spizella breweri</i> | | | |
| | Vesper Sparrow | <i>Pooecetes gramineus</i> | | | |
| | Lark Sparrow | <i>Chondestes grammacus</i> | | | |
| | Black-throated Sparrow | <i>Amphispiza bilineata</i> | | | |
| | Sage Sparrow | <i>Amphispiza belli</i> | | | |
| | Savannah Sparrow | <i>Passerculus sandwichensis</i> | | | |
| | Grasshopper Sparrow | <i>Ammodramus savannarum</i> | | | |
| | Fox Sparrow | <i>Passerella iliaca</i> | | 1 | |

| | Common Name | Scientific Name | Salmonid Relationship | Closely Associated with Riparian Wetlands | Closely Associated with Other Wetlands |
|--|-------------------------|-------------------------------|------------------------------|--|---|
| | Song Sparrow | Melospiza melodia | 1 | | |
| | Lincoln's Sparrow | Melospiza lincolnii | | 1 | |
| | White-crowned Sparrow | Zonotrichia leucophrys | | | |
| | Dark-eyed Junco | Junco hyemalis | | | |
| | Lapland Longspur | Calcarius lapponicus | | | |
| | Snow Bunting | Plectrophenax nivalis | | | |
| | Black-headed Grosbeak | Pheucticus melanocephalus | | | |
| | Lazuli Bunting | Passerina amoena | | 1 | |
| | Bobolink | Dolichonyx oryzivorus | | | |
| | Red-winged Blackbird | Agelaius phoeniceus | | | 1 |
| | Western Meadowlark | Stumella neglecta | | | |
| | Yellow-headed Blackbird | Xanthocephalus xanthocephalus | | | 1 |
| | Brewer's Blackbird | Euphagus cyanocephalus | | | |
| | Brown-headed Cowbird | Molothrus ater | | | |
| | Bullock's Oriole | Icterus bullockii | | 1 | |
| | Gray-crowned Rosy-Finch | Leucosticte tephrocotis | | | |
| | Pine Grosbeak | Pinicola enucleator | | | |
| | Purple Finch | Carpodacus purpureus | | | |
| | Cassin's Finch | Carpodacus cassinii | | | |
| | House Finch | Carpodacus mexicanus | | | |
| | Red Crossbill | Loxia curvirostra | | | |
| | White-winged Crossbill | Loxia leucoptera | | | |
| | Common Redpoll | Carduelis flammea | | | |

| | Common Name | Scientific Name | Salmonid Relationship | Closely Associated with Riparian Wetlands | Closely Associated with Other Wetlands |
|----------------|-----------------------------|----------------------------|------------------------------|--|---|
| | Pine Siskin | Carduelis pinus | | | |
| | American Goldfinch | Carduelis tristis | | | |
| | Evening Grosbeak | Coccothraustes vespertinus | | | 1 |
| | House Sparrow | Passer domesticus | | | |
| | Total Birds: | 230 | Total: 52 | 43 | 32 |
| Mammals | | | | | |
| | Masked Shrew | Sorex cinereus | 1 | | |
| | Vagrant Shrew | Sorex vagrans | 1 | | |
| | Montane Shrew | Sorex monticolus | 1 | | |
| | Water Shrew | Sorex palustris | 1 | 1 | |
| | Pacific Water Shrew | Sorex bendirii | 1 | | |
| | Trowbridge's Shrew | Sorex trowbridgii | 1 | | |
| | Merriam's Shrew | Sorex merriami | | | |
| | Shrew-mole | Neurotrichus gibbsii | | | |
| | Coast Mole | Scapanus orarius | | | |
| | California Myotis | Myotis californicus | | | |
| | Western Small-footed Myotis | Myotis ciliolabrum | | 1 | |
| | Yuma Myotis | Myotis yumanensis | | 1 | |
| | Little Brown Myotis | Myotis lucifugus | | | |
| | Long-legged Myotis | Myotis volans | | 1 | |
| | Fringed Myotis | Myotis thysanodes | | | |
| | Long-eared Myotis | Myotis evotis | | | |
| | Silver-haired Bat | Lasionycteris noctivagans | | | |
| | Western Pipistrelle | Pipistrellus hesperus | | 1 | |
| | Big Brown Bat | Eptesicus fuscus | | 1 | |
| | Hoary Bat | Lasiurus cinereus | | | |
| | Spotted Bat | Euderma maculatum | | | |
| | Townsend's Big-eared Bat | Corynorhinus townsendii | | | |

| | Common Name | Scientific Name | Salmonid Relationship | Closely Associated with Riparian Wetlands | Closely Associated with Other Wetlands |
|--|--|---------------------------------|------------------------------|--|---|
| | Pallid Bat | <i>Antrozous pallidus</i> | | 1 | |
| | American Pika | <i>Ochotona princeps</i> | | | |
| | Pygmy Rabbit | <i>Brachylagus idahoensis</i> | | | |
| | Eastern Cottontail | <i>Sylvilagus floridanus</i> | | | |
| | Nuttall's (Mountain) Cottontail | <i>Sylvilagus nuttallii</i> | | | |
| | Snowshoe Hare | <i>Lepus americanus</i> | | 1 | |
| | White-tailed Jackrabbit | <i>Lepus townsendii</i> | | | |
| | Black-tailed Jackrabbit | <i>Lepus californicus</i> | | | |
| | Mountain Beaver | <i>Aplodontia rufa</i> | | | |
| | Least Chipmunk | <i>Tamias minimus</i> | | | |
| | Yellow-pine Chipmunk | <i>Tamias amoenus</i> | | | |
| | Townsend's Chipmunk | <i>Tamias townsendii</i> | | | |
| | Yellow-bellied Marmot | <i>Marmota flaviventris</i> | | | |
| | Hoary Marmot | <i>Marmota caligata</i> | | | |
| | Townsend's Ground Squirrel | <i>Spermophilus townsendii</i> | | | |
| | Washington Ground Squirrel | <i>Spermophilus washingtoni</i> | | | |
| | Columbian Ground Squirrel | <i>Spermophilus columbianus</i> | | | |
| | Cascade Golden-mantled Ground Squirrel | <i>Spermophilus saturatus</i> | | | |
| | Eastern Fox Squirrel | <i>Sciurus niger</i> | | | |
| | Western Gray Squirrel | <i>Sciurus griseus</i> | | | |
| | Red Squirrel | <i>Tamiasciurus hudsonicus</i> | | | |
| | Douglas' Squirrel | <i>Tamiasciurus douglasii</i> | 1 | | |

| | Common Name | Scientific Name | Salmonid Relationship | Closely Associated with Riparian Wetlands | Closely Associated with Other Wetlands |
|--|----------------------------|----------------------------------|------------------------------|--|---|
| | Northern Flying Squirrel | <i>Glaucomys sabrinus</i> | 1 | | |
| | Northern Pocket Gopher | <i>Thomomys talpoides</i> | | | |
| | Great Basin Pocket Mouse | <i>Perognathus parvus</i> | | | |
| | American Beaver | <i>Castor canadensis</i> | | 1 | |
| | Western Harvest Mouse | <i>Reithrodontomys megalotis</i> | | 1 | |
| | Deer Mouse | <i>Peromyscus maniculatus</i> | 1 | 1 | |
| | Columbian Mouse | <i>Peromyscus keeni</i> | | | |
| | Northern Grasshopper Mouse | <i>Onychomys leucogaster</i> | | | |
| | Bushy-tailed Woodrat | <i>Neotoma cinerea</i> | | 1 | |
| | Southern Red-backed Vole | <i>Clethrionomys gapperi</i> | | 1 | |
| | Heather Vole | <i>Phenacomys intermedius</i> | | | |
| | Meadow Vole | <i>Microtus pennsylvanicus</i> | | 1 | |
| | Montane Vole | <i>Microtus montanus</i> | | | 1 |
| | Long-tailed Vole | <i>Microtus longicaudus</i> | | 1 | |
| | Creeping Vole | <i>Microtus oregoni</i> | | | |
| | Water Vole | <i>Microtus richardsoni</i> | | 1 | |
| | Sagebrush Vole | <i>Lemmyscus curtatus</i> | | | |
| | Muskrat | <i>Ondatra zibethicus</i> | | 1 | |
| | Northern Bog Lemming | <i>Synaptomys borealis</i> | | | 1 |
| | Black Rat | <i>Rattus rattus</i> | | | |
| | Norway Rat | <i>Rattus norvegicus</i> | | | |
| | House Mouse | <i>Mus musculus</i> | | | |
| | Western Jumping Mouse | <i>Zapus princeps</i> | | 1 | |

| | Common Name | Scientific Name | Salmonid Relationship | Closely Associated with Riparian Wetlands | Closely Associated with Other Wetlands |
|-----------------|---------------------------|------------------------|------------------------------|--|---|
| | Pacific Jumping Mouse | Zapus trinotatus | | 1 | |
| | Common Porcupine | Erethizon dorsatum | | | |
| | Nutria | Myocastor coypus | | | 1 |
| | Coyote | Canis latrans | 1 | | |
| | Red Fox | Vulpes vulpes | 1 | | |
| | Black Bear | Ursus americanus | 1 | | |
| | Raccoon | Procyon lotor | 1 | 1 | |
| | Ermine | Mustela erminea | | | |
| | Long-tailed Weasel | Mustela frenata | 1 | | |
| | Mink | Mustela vison | 1 | 1 | |
| | American Badger | Taxidea taxus | | | |
| | Striped Skunk | Mephitis mephitis | 1 | | |
| | Northern River Otter | Lutra canadensis | 1 | 1 | |
| | Mountain Lion | Puma concolor | 1 | | |
| | Bobcat | Lynx rufus | 1 | | |
| | Elk | Cervus elaphus | | | |
| | Mule Deer | Odocoileus hemionus | | | |
| | White-tailed Deer | Odocoileus virginianus | | | |
| | Bighorn Sheep | Ovis canadensis | | | |
| | Total Mammals: | 86 | Total: 19 | 22 | 3 |
| Reptiles | | | | | |
| | Painted Turtle | Chrysemys picta | | | |
| | Northern Alligator Lizard | Elgaria coerulea | | | |
| | Southern Alligator Lizard | Elgaria multicarinata | | | |
| | Short-horned Lizard | Phrynosoma douglassii | | | |
| | Sagebrush Lizard | Sceloporus graciosus | | | |

| | Common Name | Scientific Name | Salmonid Relationship | Closely Associated with Riparian Wetlands | Closely Associated with Other Wetlands |
|--|----------------------------------|-------------------------|------------------------------|--|---|
| | Western Fence Lizard | Sceloporus occidentalis | | | |
| | Side-blotched Lizard | Uta stansburiana | | | |
| | Western Skink | Eumeces skiltonianus | | | |
| | Rubber Boa | Charina bottae | | | |
| | Racer | Coluber constrictor | | | |
| | Ringneck Snake | Diadophis punctatus | | | |
| | Night Snake | Hypsiglena torquata | | | |
| | Striped Whipsnake | Masticophis taeniatus | | | |
| | Gopher Snake | Pituophis catenifer | | | |
| | Western Terrestrial Garter Snake | Thamnophis elegans | 1 | | |
| | Northwestern Garter Snake | Thamnophis ordinoides | | | |
| | Common Garter Snake | Thamnophis sirtalis | 1 | 1 | |
| | Western Rattlesnake | Crotalus viridis | | | |
| | Total Reptiles: | 18 | Total: 2 | 1 | 0 |
| | | | | | |
| | Total Species: | 349 | Total: 74 | 75 | 36 |

(IBIS 2003)

Fish species listed below are known or thought to occur within the UMM Subbasin (Duke Engineering 2001, GCPUD 2003). Status refers to listing as threatened or endangered: FE=federal endangered; FT=federal threatened; FSC=federal species of concern, FEL=Federal Emergency Listing, WC=Washington candidate. Asterisks indicate the species is non-native (introduced) to the UMM Subbasin.

Table 4 Fish species of the UMM Subbasin, WA.

| Family | Common Name/Status | Scientific Name |
|----------------------|---------------------------|---------------------------------|
| Acipenseridae | White sturgeon | <i>Acipenser transmontanus</i> |
| Salmonidae | Chinook salmon/FE,WC | <i>Oncorhynchus tshawytscha</i> |
| | Sockeye salmon | <i>Oncorhynchus nerka</i> |
| | Kokanee* | <i>Oncorhynchus nerka</i> |
| | Rainbow trout | <i>Oncorhynchus mykiss</i> |
| | Steelhead/FE,WC | <i>Oncorhynchus mykiss</i> |
| | Cutthroat trout | <i>Oncorhynchus clarki</i> |
| | Brown trout* | <i>Salmo trutta</i> |
| | Brook trout* | <i>Salvelinus fontinalis</i> |
| | Bull trout/FT,WC | <i>Salvelinus confluentus</i> |
| | Mountain whitefish | <i>Prosopium williamsoni</i> |
| | Lake whitefish | <i>Coregonis clupeaformis</i> |
| Percidae | Walleye* | <i>Stizostedion vitreum</i> |
| | Yellow perch* | <i>Perca flavescens</i> |
| Centrarcidae | Largemouth bass* | <i>Micropterus salmoides</i> |
| | Smallmouth bass* | <i>Micropterus dolomieu</i> |
| | Black crappie* | <i>Pomoxis nigromaculatus</i> |
| | White crappie* | <i>Pomoxis anularis</i> |
| | Bluegill* | <i>Lepomis macrochirus</i> |
| | Pumpkinseed* | <i>Lepomis gibbosus</i> |
| Gadidae | Burbot | <i>Lota lota</i> |
| Ictaluridae | Channel catfish* | <i>Ictalurus punctatus</i> |
| | Black bullhead* | <i>Ictalurus melas</i> |
| | Brown bullhead* | <i>Ictalurus nebulosus</i> |

| Family | Common Name/Status | Scientific Name |
|------------------------|---------------------------|----------------------------------|
| Catostomidae | Largescale sucker | <i>Catostomus macrocheilus</i> |
| | Bridgelip sucker | <i>Catostomus columbianus</i> |
| | Longnose sucker | <i>Catostomus catostomus</i> |
| | Mountain sucker/WC | <i>Catostomus platyrhynchus</i> |
| Cyprinidae | Carp* | <i>Cyprinus carpio</i> |
| | Northern pikeminnow | <i>Ptychocheilus oregonensis</i> |
| | Redside shiner | <i>Richardsonius balteatus</i> |
| | Chiselmouth | <i>Acrocheilus alutaceus</i> |
| | Peamouth | <i>Mylocheilus caurinus</i> |
| | Tench | <i>Tinca tinca</i> |
| | Longnose dace | <i>Rhinichthys cataractae</i> |
| | Speckled dace | <i>Rhinichthys osculus</i> |
| | Leopard dace | <i>Rhinichthys falcatus</i> |
| Percopsidae | Sand roller | <i>Percopsis transmontana</i> |
| Cottidae | Prickly sculpin | <i>Cottus asper</i> |
| | Torrent sculpin | <i>Cottus rhotheus</i> |
| Gasterosteidae | Threespine stickleback | <i>Gasterosteus aculeatus</i> |
| Petromyzontidae | Pacific lamprey/FSC | <i>Entosphenus tridentatus</i> |

Table 5 Threatened and endangered species of the UMM Subbasin, WA.

| | Common Name | Scientific Name | State Status | | Federal Status |
|---------------------------------|-------------------------|---------------------------|---------------------|-------------------|-----------------------|
| Amphibians | | | | | |
| | Western Toad | Bufo boreas | WA | Candidate Species | |
| | Columbia Spotted Frog | Rana luteiventris | WA | Candidate Species | |
| Total Listed Amphibians: | | 2 | | | |
| Birds | | | | | |
| | Common Loon | Gavia immer | WA | Sensitive | |
| | Western Grebe | Aechmophorus occidentalis | WA | Candidate Species | |
| | Northern Goshawk | Accipiter gentilis | WA | Candidate Species | |
| | Ferruginous Hawk | Buteo regalis | WA | Threatened | |
| | Golden Eagle | Aquila chrysaetos | WA | Candidate Species | |
| | Bald Eagle | | WA | Threatened | Threatened |
| | Sage Grouse | Centrocercus urophasianus | WA | Threatened | Candidate Species |
| | Sharp-tailed Grouse | Tympanuchus phasianellus | WA | Threatened | |
| | Burrowing Owl | Athene cunicularia | WA | Candidate Species | |
| | Spotted Owl | Strix occidentalis | WA | Endangered | Threatened |
| | Vaux's Swift | Chaetura vauxi | WA | Candidate Species | |
| | Lewis' Woodpecker | Melanerpes lewis | WA | Candidate Species | |
| | White-headed Woodpecker | Picoides albolarvatus | WA | Candidate Species | |
| | Black-backed Woodpecker | Picoides arcticus | WA | Candidate Species | |

| | Common Name | Scientific Name | State Status | | Federal Status |
|-------------------------------|----------------------------|--------------------------|---------------------|-------------------|-----------------------|
| | Pileated Woodpecker | Dryocopus pileatus | WA | Candidate Species | |
| | Loggerhead Shrike | Lanius ludovicianus | WA | Candidate Species | |
| | White-breasted Nuthatch | Sitta carolinensis | WA | Candidate Species | |
| | Sage Thrasher | Oreoscoptes montanus | WA | Candidate Species | |
| | Vesper Sparrow | Poocetes gramineus | WA | Candidate Species | |
| | Sage Sparrow | Amphispiza belli | WA | Candidate Species | |
| Total Listed Birds: | | 19 | | | |
| Mammals | | | | | |
| | Merriam's Shrew | Sorex merriami | WA | Candidate Species | |
| | Townsend's Big-eared Bat | Corynorhinus townsendii | WA | Candidate Species | |
| | Pygmy Rabbit | Brachylagus idahoensis | WA | Endangered | Endangered |
| | White-tailed Jackrabbit | Lepus townsendii | WA | Candidate Species | |
| | Black-tailed Jackrabbit | Lepus californicus | WA | Candidate Species | |
| | Washington Ground Squirrel | Spermophilus washingtoni | WA | Candidate Species | Candidate Species |
| | Western Gray Squirrel | Sciurus griseus | WA | Threatened | |
| | Northern Pocket Gopher | Thomomys talpoides | WA | Candidate Species | |
| Total Listed Mammals: | | 8 | | | |
| Reptiles | | | | | |
| | Striped Whipsnake | Masticophis taeniatus | WA | Candidate Species | |
| Total Listed Reptiles: | | 1 | | | |
| | | | | | |
| Total Listed Species: | | 30 | | | |

(IBIS 2003)

Table 6 Partners in Flight species of the UMM Subbasin, WA.

| Common Name | Scientific Name | PIF 1998-1999 Continental | PIF Ranking by Super Region Draft 2002 | WA PIF Priority & Focal Species |
|------------------------|----------------------------------|----------------------------------|---|--|
| Northern Harrier | <i>Circus cyaneus</i> | | | Yes |
| Swainson's Hawk | <i>Buteo swainsoni</i> | | MO (Intermountain West, Prairies) | Yes |
| Ferruginous Hawk | <i>Buteo regalis</i> | | | Yes |
| Rough-legged Hawk | <i>Buteo lagopus</i> | | PR (Arctic) | |
| American Kestrel | <i>Falco sparverius</i> | | | Yes |
| Gyr Falcon | <i>Falco rusticolus</i> | | PR (Arctic) | |
| Sage Grouse | <i>Centrocercus urophasianus</i> | | MA (Intermountain West, Prairies) | |
| Spruce Grouse | <i>Falcapennis canadensis</i> | | PR (Northern Forests) | |
| White-tailed Ptarmigan | <i>Lagopus leucurus</i> | | MO (Arctic) | |
| Blue Grouse | <i>Dendragapus obscurus</i> | | MA (Pacific, Intermountain West) | |
| Sharp-tailed Grouse | <i>Tympanuchus phasianellus</i> | | MO (Prairies) | Yes |
| Long-billed Curlew | <i>Numenius americanus</i> | Yes | | |
| Stilt Sandpiper | <i>Calidris himantopus</i> | Yes | | |
| Flammulated Owl | <i>Otus flammeolus</i> | | MO (Pacific, Intermountain West, Southwest) | Yes |
| Snowy Owl | <i>Nyctea scandiaca</i> | | PR (Arctic) | |
| Northern Pygmy-owl | <i>Glaucidium gnoma</i> | | PR (Pacific) | |
| Burrowing Owl | <i>Athene cunicularia</i> | | | Yes |
| Spotted Owl | <i>Strix occidentalis</i> | | IM (Pacific, Intermountain West, Southwest) | |
| Great Gray Owl | <i>Strix nebulosa</i> | | | Yes |
| Short-eared Owl | <i>Asio flammeus</i> | Yes | MA (Arctic, Northern Forests, Intermountain West, Prairies) | Yes |
| Common Poorwill | <i>Phalaenoptilus nuttallii</i> | | | Yes |
| Black Swift | <i>Cypseloides niger</i> | Yes | IM (Pacific, Intermountain) | Yes |

| Common Name | Scientific Name | PIF 1998-1999 Continental | PIF Ranking by Super Region Draft 2002 | WA PIF Priority & Focal Species |
|-------------------------|-------------------------------|----------------------------------|--|--|
| | | | West) | |
| Vaux's Swift | <i>Chaetura vauxi</i> | | | Yes |
| Calliope Hummingbird | <i>Stellula calliope</i> | | MO (Intermountain West) | Yes |
| Rufous Hummingbird | <i>Selasphorus rufus</i> | Yes | MA (Pacific, Intermountain West) | Yes |
| Lewis's Woodpecker | <i>Melanerpes lewis</i> | Yes | MO (Intermountain West, Prairies) | Yes |
| Williamson's Sapsucker | <i>Sphyrapicus thyroideus</i> | | MO (Intermountain West) | Yes |
| Red-naped Sapsucker | <i>Sphyrapicus nuchalis</i> | | MO (Intermountain West) | Yes |
| Red-breasted Sapsucker | <i>Sphyrapicus ruber</i> | | MO (Pacific) | Yes |
| Downy Woodpecker | <i>Picoides pubescens</i> | | | Yes |
| White-headed Woodpecker | <i>Picoides albolarvatus</i> | Yes | PR (Pacific, Intermountain West) | Yes |
| Three-toed Woodpecker | <i>Picoides tridactylus</i> | | PR (Northern Forests) | |
| Black-backed Woodpecker | <i>Picoides arcticus</i> | | PR (Northern Forests) | Yes |
| Pileated Woodpecker | <i>Dryocopus pileatus</i> | | | Yes |
| Olive-sided Flycatcher | <i>Contopus cooperi</i> | | MA (Pacific, Northern Forests, Intermountain West) | Yes |
| Western Wood-pewee | <i>Contopus sordidulus</i> | | | Yes |
| Willow Flycatcher | <i>Empidonax traillii</i> | | MA (Prairies, East) | Yes |
| Hammond's Flycatcher | <i>Empidonax hammondii</i> | | | Yes |
| Gray Flycatcher | <i>Empidonax wrightii</i> | | PR (Intermountain West) | Yes |
| Dusky Flycatcher | <i>Empidonax oberholseri</i> | | MA (Intermountain West) | Yes |

| Common Name | Scientific Name | PIF 1998-1999 Continental | PIF Ranking by Super Region Draft 2002 | WA PIF Priority & Focal Species |
|---------------------------|--------------------------------|----------------------------------|---|--|
| Pacific-slope Flycatcher | <i>Empidonax difficilis</i> | | PR (Pacific) | Yes |
| Loggerhead Shrike | <i>Lanius ludovicianus</i> | | | Yes |
| Northern Shrike | <i>Lanius excubitor</i> | | PR (Northern Forests) | |
| Warbling Vireo | <i>Vireo gilvus</i> | | | Yes |
| Red-eyed Vireo | <i>Vireo olivaceus</i> | | | Yes |
| Gray Jay | <i>Perisoreus canadensis</i> | | PR (Northern Forests) | |
| Clark's Nutcracker | <i>Nucifraga columbiana</i> | | PR (Intermountain West) | Yes |
| Horned Lark | <i>Eremophila alpestris</i> | | | Yes |
| Bank Swallow | <i>Riparia riparia</i> | | | Yes |
| Chestnut-backed Chickadee | <i>Poecile rufescens</i> | | PR (Pacific) | |
| Boreal Chickadee | <i>Poecile hudsonicus</i> | | MA (Northern Forests) | |
| White-breasted Nuthatch | <i>Sitta carolinensis</i> | | | Yes |
| Brown Creeper | <i>Certhia americana</i> | | | Yes |
| House Wren | <i>Troglodytes aedon</i> | | | Yes |
| Winter Wren | <i>Troglodytes troglodytes</i> | | | Yes |
| American Dipper | <i>Cinclus mexicanus</i> | | | Yes |
| Western Bluebird | <i>Sialia mexicana</i> | | | Yes |
| Mountain Bluebird | <i>Sialia currucoides</i> | | PR (Intermountain West) | |
| Townsend's Solitaire | <i>Myadestes townsendi</i> | | | Yes |
| Veery | <i>Catharus fuscescens</i> | | | Yes |
| Swainson's Thrush | <i>Catharus ustulatus</i> | | | Yes |
| Hermit Thrush | <i>Catharus guttatus</i> | | | Yes |
| Varied Thrush | <i>Ixoreus naevius</i> | | | Yes |
| Sage Thrasher | <i>Oreoscoptes montanus</i> | | PR (Intermountain West) | Yes |
| American Pipit | <i>Anthus rubescens</i> | | PR (Arctic) | Yes |
| Bohemian Waxwing | <i>Bombycilla garrulus</i> | | MA (Northern | |

| Common Name | Scientific Name | PIF 1998-1999 Continental | PIF Ranking by Super Region Draft 2002 | WA PIF Priority & Focal Species |
|-----------------------------|----------------------------------|---------------------------|--|---------------------------------|
| | | | Forests) | |
| Orange-crowned Warbler | <i>Vermivora celata</i> | | | Yes |
| Nashville Warbler | <i>Vermivora ruficapilla</i> | | PR (Northern Forests) | Yes |
| Yellow Warbler | <i>Dendroica petechia</i> | | | Yes |
| Yellow-rumped Warbler | <i>Dendroica coronata</i> | | | Yes |
| Black-throated Gray Warbler | <i>Dendroica nigrescens</i> | | MO (Pacific) | Yes |
| Townsend's Warbler | <i>Dendroica townsendi</i> | | | Yes |
| Hermit Warbler | <i>Dendroica occidentalis</i> | Yes | MO (Pacific) | Yes |
| Macgillivray's Warbler | <i>Oporornis tolmiei</i> | | | Yes |
| Wilson's Warbler | <i>Wilsonia pusilla</i> | | | Yes |
| Yellow-breasted Chat | <i>Icteria virens</i> | | | Yes |
| Western Tanager | <i>Piranga ludoviciana</i> | | | Yes |
| Chipping Sparrow | <i>Spizella passerina</i> | | | Yes |
| Brewer's Sparrow | <i>Spizella breweri</i> | Yes | MA (Intermountain West) | Yes |
| Vesper Sparrow | <i>Pooecetes gramineus</i> | | | Yes |
| Lark Sparrow | <i>Chondestes grammacus</i> | | | Yes |
| Black-throated Sparrow | <i>Amphispiza bilineata</i> | | | Yes |
| Sage Sparrow | <i>Amphispiza belli</i> | Yes | PR (Intermountain West) | Yes |
| Fox Sparrow | <i>Passerella iliaca</i> | | | Yes |
| Lincoln's Sparrow | <i>Melospiza lincolni</i> | | PR (Northern Forests) | Yes |
| Lapland Longspur | <i>Calcarius lapponicus</i> | | PR (Arctic) | |
| Snow Bunting | <i>Plectrophenax nivalis</i> | | PR (Arctic) | |
| Black-headed Grosbeak | <i>Pheucticus melanocephalus</i> | | | Yes |
| Bobolink | <i>Dolichonyx oryzivorus</i> | Yes | | |
| Western Meadowlark | <i>Sturnella neglecta</i> | | | Yes |
| Bullock's Oriole | <i>Icterus bullockii</i> | | | Yes |
| Pine Grosbeak | <i>Pinicola enucleator</i> | | MO (Northern Forests) | |

| Common Name | Scientific Name | PIF 1998-1999 Continental | PIF Ranking by Super Region Draft 2002 | WA PIF Priority & Focal Species |
|------------------------|-----------------------------|----------------------------------|---|--|
| Purple Finch | <i>Carpodacus purpureus</i> | | | Yes |
| Cassin's Finch | <i>Carpodacus cassinii</i> | | MA (Intermountain West) | |
| Red Crossbill | <i>Loxia curvirostra</i> | | | Yes |
| White-winged Crossbill | <i>Loxia leucoptera</i> | | PR (Northern Forests) | |
| Total Species: | 98 | | | |

(IBIS 2003)

Table 7 Wildlife game species of the UMM Subbasin, WA.

| | Common Name | Scientific Name | WA |
|-------------------|-------------------------------|-------------------------------|--------------|
| Amphibians | | | |
| | Bullfrog | <i>Rana catesbeiana</i> | Game Species |
| | Total Game Amphibians: | 1 | |
| Birds | | | |
| | Canada Goose | <i>Branta canadensis</i> | Game Bird |
| | Wood Duck | <i>Aix sponsa</i> | Game Bird |
| | Gadwall | <i>Anas strepera</i> | Game Bird |
| | American Wigeon | <i>Anas americana</i> | Game Bird |
| | Mallard | <i>Anas platyrhynchos</i> | Game Bird |
| | Blue-winged Teal | <i>Anas discors</i> | Game Bird |
| | Cinnamon Teal | <i>Anas cyanoptera</i> | Game Bird |
| | Northern Shoveler | <i>Anas clypeata</i> | Game Bird |
| | Northern Pintail | <i>Anas acuta</i> | Game Bird |
| | Green-winged Teal | <i>Anas crecca</i> | Game Bird |
| | Canvasback | <i>Aythya valisineria</i> | Game Bird |
| | Redhead | <i>Aythya americana</i> | Game Bird |
| | Ring-necked Duck | <i>Aythya collaris</i> | Game Bird |
| | Greater Scaup | <i>Aythya marila</i> | Game Bird |
| | Barrow's Goldeneye | <i>Bucephala islandica</i> | Game Bird |
| | Hooded Merganser | <i>Lophodytes cucullatus</i> | Game Bird |
| | Common Merganser | <i>Mergus merganser</i> | Game Bird |
| | Ruddy Duck | <i>Oxyura jamaicensis</i> | Game Bird |
| | Chukar | <i>Alectoris chukar</i> | Game Bird |
| | Gray Partridge | <i>Perdix perdix</i> | Game Bird |
| | Ring-necked Pheasant | <i>Phasianus colchicus</i> | Game Bird |
| | Ruffed Grouse | <i>Bonasa umbellus</i> | Game Bird |
| | Spruce Grouse | <i>Falciennis canadensis</i> | Game Bird |
| | Blue Grouse | <i>Dendragapus obscurus</i> | Game Bird |
| | Wild Turkey | <i>Meleagris gallopavo</i> | Game Bird |
| | California Quail | <i>Callipepla californica</i> | Game Bird |

| | Common Name | Scientific Name | WA |
|----------------|---------------------------------|------------------------|-------------|
| | American Coot | Fulica americana | Game Bird |
| | Common Snipe | Gallinago gallinago | Game Bird |
| | Mourning Dove | Zenaida macroura | Game Bird |
| | Total Game Birds: | 29 | |
| Mammals | | | |
| | Eastern Cottontail | Sylvilagus floridanus | Game Mammal |
| | Nuttall's (Mountain) Cottontail | Sylvilagus nuttallii | Game Mammal |
| | Snowshoe Hare | Lepus americanus | Game Mammal |
| | White-tailed Jackrabbit | Lepus townsendii | Game Mammal |
| | Black-tailed Jackrabbit | Lepus californicus | Game Mammal |
| | American Beaver | Castor canadensis | Game Mammal |
| | Muskrat | Ondatra zibethicus | Game Mammal |
| | Red Fox | Vulpes vulpes | Game Mammal |
| | Black Bear | Ursus americanus | Game Mammal |
| | Raccoon | Procyon lotor | Game Mammal |
| | Ermine | Mustela erminea | Game Mammal |
| | Long-tailed Weasel | Mustela frenata | Game Mammal |
| | Mink | Mustela vison | Game Mammal |
| | American Badger | Taxidea taxus | Game Mammal |
| | Northern River Otter | Lutra canadensis | Game Mammal |
| | Mountain Lion | Puma concolor | Game Mammal |
| | Bobcat | Lynx rufus | Game Mammal |
| | Rocky Mountain Elk | Cervus elaphus nelsoni | Game |

| | Common Name | Scientific Name | WA |
|--|----------------------------|------------------------|----------------|
| | | | Mammal |
| | Mule Deer | Odocoileus hemionus | Game Mammal |
| | Bighorn Sheep | Ovis canadensis | Game Mammal |
| | Total Game Mammals: | 20 | |
| | | | |
| | Total Game Species: | 50 | |

(IBIS 2003).

Appendix C

Upper Middle Mainstem Columbia River Subbasin

Focal Species Information, Red-winged Blackbird

Introduction

The red-winged black bird is one of the most abundant birds in North America (Marshall et al. 2003). Red-winged Blackbirds are extremely adaptable; successfully colonizing many small wetlands created by human activities (i.e., farming, road building, and industrial wetlands) Loss of natural wetlands has frequently been compensated for by these human activities (UW 1991). The bird is considered a pest species in many areas where huge flocks damage crops.

Life History

Diet

About 75% of the annual Red-winged Blackbird diet is seeds. During the breeding season, they also eat insects, especially dragonflies, mayflies, and caddis flies as they emerge from their aquatic larval stage. In winter, grain is an important source of food, and many birds feed on corn stubble and at feedlots (SAS 2002). The species sometimes forms large, sexually separate flocks in wetland herbaceous habitats, trees, brushlands, and feedlots, and may forage on agricultural crops (i.e., corn, rice, oat, wheat, alfalfa, and sunflower) or on understory seed sources (Mott et al. 1972; Johnson and Caslick 1982, Marshall et al. 2003). During nesting season, red-winged blackbirds may forage within the understory, midstory, and overstory canopies of the wetland they are nesting in, or within a nearby wetland (Snelling 1968, Holm 1973).

Reproduction

The timing of breeding varies throughout the range of the red-winged blackbird. Nesting frequently begins in March or April and is completed by mid-July in the more temperate habitats (Short 1985). Older males (2+ yr) return to breeding sites first, followed by adult females and younger birds. Females nest as yearlings, males not until the second year (Marshall et al. 2003). Males are highly territorial and polygynous; up to six females commonly nest within a male's territory. Females sometimes mate with several partners during a season or even during a single nesting attempt. Males do not participate in nest building, incubation, or feeding of the incubating female (G.H. Orians, pers. comm., 1984). Males sometimes feed older nestlings and fledglings (Marshall et al. 2003). Most young in North America are fledged by late July (Short 1985).

Nesting

The red-winged blackbird nests in fresh-water and brackish herbaceous wetlands, shrubs (Douglas spiraea, small Oregon ash, willow, and alder trees) and small trees (i.e.,

willows) along watercourses (AOU 1983:723, Marshall et al. 2003), in upland habitat (grass, forb, and pasture/hay cover types, roadsides, canals, ditches and parks and suburban habitat) near surface water, and in suitable vegetation distant from free water (Dolbeer 1980, Micacchion and Townsend 1983, Marshall et al. 2003). Herbaceous wetlands or sloughs, with extensive cattails, bulrushes, sedges, reeds (*Phragmites spp.*), or tules (*Scirpus spp.*), historically have provided important nesting habitat for the blackbird (Bent 1958).

Females select the nest sites and build the nests. They are made of grass and are usually lashed to cattails, bulrushes, or other emergent vegetation about 8-32 in (0.2-0.8 m) above water (Marshall et al. 2003). Red-winged blackbirds seem to prefer areas with the densest, tallest herbaceous vegetation for nest placement (trees greater than 5.0 m in height) (Albers 1978). Nests that border areas of open water are placed on the edges of cattail clumps (Wiens 1965), while those in upland sites typically are wound between and attached to stalks of herbaceous vegetation (Bent 1958). Herbaceous wetlands that are dominated by cattails and have open, permanent water have the optimum number of available nest sites (Weatherhead and Robertson 1977). Early nests are placed in robust, dense, old herbaceous growth and are more productive than late nests, which are entwined with stems of the new growth (Meanley and Webb 1963).

Red-winged blackbirds may lay as many as 5 eggs, but usually 3-4. Young fledge 12 days after hatching. Parents feed fledglings for 30 days after fledging (Marshall et al. 2003).

Nest success seems to be related to presence or absence of permanent water, water depth (greater nest success in water up to 50 cm or more) within the wetland, proximity of the nest to water (greater for nests 20 cm above water than those 100 cm above water), relative openness of nesting cover within the wetland, and the type of vegetation holding the nest. Nests placed in herbaceous wetland vegetation fared better when placed where open water, marsh grass and loosestrife (vs. sweet gale and sedges) were present (Weatherhead and Robertson 1977).

Herbaceous wetlands dominated by cattails generally seem to be the most productive habitats for red-winged blackbirds in terms of nests/ha or number of young fledged/ha (Robertson 1972). Favorable herbaceous wetland sites produce more suitable food per unit area and have higher nest densities, highly synchronous nesting, higher nest survival rates, and lower nest predation rates than do upland nest sites (Short 1985).

Migration

Some populations in the southern parts of the range are nonmigratory, but almost all northern birds winter in the South, forming huge flocks that migrate by day, foraging for grain and seeds in fields with other blackbirds, and roost at night in dense cover in wetland habitats (SAS, 2002). Males migrate to or congregate at future nesting habitats in late winter, and females arrive at the territories in early spring (Case and Hewitt 1963). In areas with resident populations, individuals of both sexes may remain near breeding territories throughout the year, even though the areas are not actively defended or used in winter except, perhaps, as roosting sites (G.H. Orians, pers. comm., 1984).

Mortality

Marsh wrens peck at red-winged blackbird eggs and the northern harrier, American crow, and raccoon predate the nests. Nesting success increases with nest dispersion and distance from marsh wrens. Nearby nesting females also reduces predation risk (Marshall et al. 2003), and the presence of permanent water within the wetland may reduce mammalian predation on nests (Robertson 1972).

In addition, the abundance of red-winged blackbirds is negatively correlated with the presence and abundance of carp, along with disturbances such as grazing, mowing, burning, and tilling of potential upland nest sites. Carp disturb submerged wetland vegetation and destroy food sources (aquatic insects) for the blackbird. Activities such as grazing and mowing destroy potential nesting habitat and interfere with nesting birds.

Habitat Requirements

Red-winged blackbirds need tall, dense, persistent herbaceous vegetation reasonably close to water for nesting, foraging, and cover requirements, whether it be in a wetland or upland environment. The bird readily uses midstory and overstory layers of habitat at times but does not seem to be dependent on the presence of these layers (Short 1985). In a wetland environment, blackbirds prefer patchy stands of cattails interspersed with areas of open water, over dense homogeneous stands of cattails (Robertson 1972). An important characteristic of upland nest sites is the availability of fence posts and other structures that serve as display perches for males and as observation posts for both males and females (Joyner 1978).

Blackbirds also require an abundant supply of aquatic insects for foraging in the spring and early summer. Wetlands that are permanently flooded, or intermittently exposed, with water usually present throughout the year are necessary to support persistent populations of submergent vegetation and benthic invertebrates (Orians 1980).

The red-winged blackbird does not require large territories and are often seen in very small patches of habitat (SAS 2002). In winter they often congregate in agricultural areas. Short (1985) surmised that a wetland area must contain at least 0.10 ha in emergent herbaceous vegetation, like cattails, to be considered nesting habitat for the blackbird. Several studies have described the minimum territory for male red-winged blackbirds as 0.02 ha (Weatherhead and Robertson 1977; Orians 1980). Territories in upland habitats are much larger, requiring at least 1.0 ha in area to provide adequate breeding habitat for the bird (Short 1985).

Focal Species Population and Distribution

Population

Current

The red-winged blackbird is one of the most abundant species of bird in North America, with an estimated 190 million-winter population. The red-winged blackbird breeds from southeast Alaska across Canada to south central Quebec, and south to the Caribbean, Mexico, and Middle America. It winters from southeast Alaska and Canada, south to the Gulf Coast and Mexico. It is also a widespread and abundant breeder throughout

Washington's lowlands. There are sixteen subspecies in North America (Marshall et al. 2003) and two poorly distinguished subspecies in Washington: *A.p. caurinus* of western Washington and *A.p. nevadensis* of eastern Washington (UW 1991).

Distribution

Current

This aggressive species is widespread and abundant at lower elevations of the State of Washington, including the UMM Subbasin, in virtually every habitat as long as a suitable microhabitat with emergent vegetation is available. It can also be found along roads where ditches have created suitable habitat. They rarely nest in upland shrubby areas (UW 1991). In winter they are often less widespread, but can be found year round on the Columbia River (BirdWeb 2003)

In a University of Washington study, core areas of habitat were all water / wetlands (including estuaries) below the subalpine fir zone. All other habitats except bare ground were suitable if small pockets of wetland occurred within the larger mapped habitats (UW 1991).

The red-winged blackbird is distributed throughout the UMM Subbasin with confirmed, probable and possible breeding sites. Confirmed sitings are primarily along the Columbia River (near cities of Vantage, Wenatchee, Pateros, Bridgeport and Rocky Butte), Banks Lake, and Lake Lenore State Wildlife Recreation Area (WDFW 1999). Nesting red-winged black birds are abundant on herbaceous wetlands in the northern portion of the UMM (Braaten, pers. comm., 2004).

Focal Species Status and Abundance Trends

Status

The red-winged blackbird is one of the most abundant species in North America with an estimated winter population of 190 million (Marshall et al. 2003). This species is also a common summer resident in the wetlands and marshes of Washington State and is a common winter visitor on farmlands (SAS 2002). In the Dakotas, redwings have declined because of drought and tilling of breeding areas (Marshall et al. 2003). This species is not currently listed as endangered or threatened by the federal or state government.

The blackbird is highly efficient in adapting to anthropogenic environments and has had a significant impact on agricultural crops. Winter roosts can be huge, especially in major grain-producing areas like Washington. Costs related to their consumption of grain (wheat, barley, corn, sunflower and rice) can become high and may exceed the benefits of insect control related to their foraging habits during fledging (Bendell et al. 1981). Grain fields closest to blackbird roost areas have comparably greater economic losses. Poisoning, trapping, shooting, or flock harassment by loud noises is allowed by an amendment to the Federal Migratory Bird Treaty Act of 1918. Other means of population control include reducing grain waste, using resistant cultivars and crops less favorable to blackbirds, and timing of agricultural activities (Marshall et al. 2003).

Trends

The North American BBS trend estimates for the red-winged blackbird within the state of Washington are .5% (1966-2002), -2.2% (1996-1979), and .1% (1980-2002) change in population per year. BBS data indicate there is a less than 1.5% decreasing trend (1966-1996) in red-winged blackbird populations within the UMM (Sauer et al. 2003).

Key Factors Inhibiting Populations and Ecological Processes

Activities, such as intensive livestock grazing, mowing, burning, and tilling of old growth stubble, make herbaceous uplands unavailable for early nest placement. Mowing hayfields during the nesting season disrupts nesting success on upland sites (Albers 1978).

The presence and abundance of carp within a wetland may inhibit red-winged blackbird populations. Carp disturb submergent vegetation within the wetland, which may destroy habitat for emergent aquatic insects (like Odonates) and reduce food sources for blackbirds (Short 1985).

Albers, P.H. 1978. Habitat selection by breeding red-winged blackbirds. *Wilson Bulletin*, 90(4):619-634.

American Ornithologists' Union. 1983 checklist of North American birds. 6th edition. Am. Ornithologists Union. 877 pp.

Bendell, B., P. Weatherhead, and R. Stewart. 1981. The impact of predation by red-winged blackbirds on European corn borer populations. *Canadian Journal of Zoology* 59(8): 1535-1538.

Bent, A. C. 1958. Life histories of North American blackbirds, orioles, tanagers, and allies. U.S. National Museum Bulletin, 211. 549 pp.

BirdWeb. 2003. Birds of Washington maps: red-winged blackbird distribution in Washington State. http://users.pullman.com/lostriver/sas_maps.html. Feb. 2004.

Braaten, E. 2004. Wildlife biologist, Washington Department of Fish and Wildlife, Electric City, WA.

Case, N.S., and O.H. Hewitt. 1963. Nesting and productivity of the re-winged blackbird in relation to habitat. *Living Bird* 2:7-20.

Dolbeer, R.A. 1980. Blackbirds and corn in Ohio. U.S. fish and Wildlife Service. Res. Publication #136. 18pp.

Holm, C.H. 1973. Breeding sex ratios, territoriality, and reproductive success in the red-winged blackbird (*Agelaius phoeniceus*). *Ecology* 54(2):356-365.

Johnson, R.J., and J.W. Caslick. 1982. Habitat relationships of roosting and flocking red-winged blackbirds. *Journal of Wildlife Management*, 46(4):1071-1077.

Joyner, D.E. 1978. Use of an old-field habitat by boblinks and red-winged blackbirds. *Canadian Field-Naturalist* 92(4):383-386.

Marshall, D., M. Hunter, and A. Contreras, Eds. 2003. *Birds of Oregon: A general reference*. Oregon State University Press, Corvallis, OR. 768 pp.

Meanley, B. and J. Webb. 1963. Nesting ecology and reproductive rate of the red-winged blackbird in tidal marshes of the upper Chesapeake Bay region. *Chesapeake Science* 4(2):90-100.

- Micacchion, M., and T.W. Townsend. 1983. Botanical characteristics of autumnal blackbird roosts in central Ohio. *Ohio Academy of Science*, 83(3):131-135.
- Mott, D.F., R. R. West, J.W. DeGrazio, and J.L. Guarino. 1972. Foods of the red-winged blackbird in Brown County, South Dakota. *Journal of Wildlife Management* 36(3):983-987.
- Orians, G.H. 1980. Some adaptations of marsh-nesting blackbirds. Princeton University press, Princeton, NJ. 295 pp.
- . Personal communication (letter dated August 14, 1984) with H. L. Short. University of Washington, Seattle, WA. Citation in Short, H.L. 1985. Habitat suitability index models: Red-winged blackbird. U.S. Fish and Wildlife Service, Biological Report 82(10.95). 20 pp.
- Robertson, R.J. 1972. Optimal niche space of the red-winged blackbird (*Agelaius phoeniceus*). Nesting success in marsh and upland habitat.
- Sauer, J. R., J. E. Hines, and J. Fallon. 2003. The North American Breeding Bird Survey, Results and Analysis 1966 - 2002. Version 2003.1, USGS Patuxent Wildlife Research Center, Laurel, MD.
- Seattle Audubon Society (SAS). 2002. BirdWeb: red-winged blackbird. <http://www.birdweb.org/birdweb/Species.asp?id=438>. Feb. 2004.
- Short, H.L. 1985. Habitat suitability index models: Red-winged blackbird. U.S. Fish and Wildlife Service, Biological Report 82(10.95). 20 pp.
- Snelling, J.C. 1968. Overlap in feeding habitats of red-winged blackbirds and common grackles nesting in a cattail marsh. *Auk* 85(4):560-585.
- University of Washington (UW). 1991. Nature mapping, Washington gap analysis projects predicted distribution Map, red-winged blackbird. <http://www.fish.washington.edu/naturemapping/maphtml/bagph.html>. Feb. 2004.
- Washington Department of Fish and Wildlife (WDFW). 1999. Washington gap data products: red-winged blackbird. <http://WDFW.wa.gov/wlm/gap/birds.htm>. Feb. 2004
- Weatherhead, P.J., and R.J. Robertson. 1977. Harem size, territory quality, and reproductive success in the red-winged blackbird (*Agelaius phoeniceus*). *Canadian Journal of Zoology*, 55(8):1261-1267.
- Wiens, J.A. 1965. Behavioral interactions of red-winged blackbirds and common grackles on a common breeding ground. *Auk* 82(3):356-374.

Appendix D

Upper Middle Mainstem Columbia River Subbasin

Conservation Reserve Program

Table 1 Conservation Reserve Program acreage, UMM Subbasin, WA.

| Washington Report ID - MEPRTN-R1 | | U.S. Department of Agriculture - Farm Service Agency Conservation Reserve Program - Monthly Contract Report Summary for Active Contracts for all Program Years (1986-2004) | | | | | As of: 08-29-03 Prepared on: 08-29-03 Page: 13 | | | |
|--|---------------------------|--|---------------------|--------------------------|---------------------------------|-----------------------------|--|------------------------|-----------------------|--|
| County Name | Total No. of Contracts | Total CRP Acres | Avg. Rental Rate | Continuous CREP Acres | Continuous Non-Crep Acres | Wetland Systems Acres | Margin Pastureland Acres | Tree Practice Acres | Avg. Erosion Index | |
| ADAMS | 1,696 | 212,463.9 | \$50.17 | 0 | 17,206.1 | 207.0 | 0.0 | 54.0 | 5 | |
| ASOTIN | 144 | 29,145.6 | \$54.28 | 760.5 | 111.6 | 0.0 | 852.3 | 907.1 | 11 | |
| BENTON | 402 | 74,265.9 | \$39.93 | 0 | 5,896.3 | 0.0 | 0.0 | 0.0 | 9 | |
| CHELAN | 8 | 1,372.7 | \$47.01 | 4.5 | 0.0 | 0.0 | 0.0 | 4.5 | 6 | |
| CLALLAM | 6 | 34.3 | \$159.03 | 34.3 | 0.0 | 0.0 | 33.3 | 34.3 | 3 | |
| CLARK | 7 | 76.7 | \$145.65 | 62.3 | 14.4 | 0.0 | 76.7 | 76.7 | 76 | |
| COLUMBIA | 306 | 38,583.8 | \$61.87 | 1,424.90 | 507.1 | 0.0 | 1,714.0 | 2,841.2 | 15 | |
| COWLITZ | 2 | 14.8 | \$163.96 | 14.8 | 0.0 | 0.0 | 14.8 | 14.8 | 1 | |
| DOUGLAS | 1,076 | 187,711.0 | \$45.36 | 0 | 747.5 | 533.7 | 60.5 | 150.0 | 5 | |
| FERRY | 17 | 1,090.7 | \$55.01 | 0 | 25.4 | 0.0 | 0.0 | 14.5 | 13 | |
| FRANKLIN | 776 | 104,426.7 | \$50.35 | 0 | 12,727.8 | 0.0 | 4.6 | 8.1 | 5 | |
| GARFIELD | 464 | 44,655.1 | \$65.80 | 650.9 | 2,493.9 | 89.9 | 2,027.8 | 2,225.2 | 14 | |
| GRANT | 405 | 60,715.5 | \$43.85 | 0 | 1,117.8 | 0.0 | 0.0 | 0.0 | 7 | |
| GRAYS HARBOR | 11 | 105.0 | \$183.46 | 74.7 | 30.3 | 0.0 | 77.2 | 105.0 | 1 | |
| JEFFERSON | 9 | 97.2 | \$220.10 | 97.2 | 0.0 | 0.0 | 76.5 | 97.2 | 15 | |
| KING | 1 | 5.3 | \$204.40 | 5.3 | 0.0 | 0.0 | 5.3 | 5.3 | 1 | |
| KITSAP | 1 | 5.0 | \$199.60 | 5 | 0.0 | 0.0 | 5.0 | 5.0 | 243 | |
| KITTITAS | 19 | 3,294.2 | \$50.62 | 0 | 0.0 | 0.0 | 0.0 | 0.0 | 18 | |
| KLICKITAT | 360 | 58,407.9 | \$44.03 | 47.5 | 4,598.3 | 0.0 | 4,130.4 | 4,378.0 | 9 | |
| LEWIS | 24 | 515.8 | \$188.17 | 436.4 | 79.4 | 0.0 | 449.5 | 498.9 | 1 | |
| LINCOLN | 955 | 86,270.7 | \$46.18 | 0 | 1,644.1 | 857.7 | 16.9 | 388.4 | 8 | |
| MASON | 6 | 37.3 | \$191.68 | 37.3 | 0.0 | 0.0 | 37.3 | 37.3 | 1 | |
| OKANOGAN | 50 | 4,064.6 | \$49.11 | 33.9 | 50.0 | 2,737.3 | 0.0 | 83.9 | 4 | |
| PACIFIC | 3 | 41.4 | \$211.16 | 41.4 | 0.0 | 0.0 | 41.4 | 41.4 | 1 | |
| PIERCE | 4 | 18.5 | \$164.94 | 3 | 15.5 | 0.0 | 5.5 | 18.5 | 10 | |
| SKAGIT | 66 | 443.4 | \$268.69 | 443.4 | 0.0 | 0.0 | 203.8 | 443.4 | 2 | |
| SNOHOMISH | 12 | 135.6 | \$229.49 | 111.8 | 23.8 | 0.0 | 127.0 | 135.6 | 5 | |
| SPOKANE | 459 | 31,768.2 | \$56.76 | 0 | 758.2 | 2,239.6 | 268.6 | 746.0 | 11 | |
| STEVENS | 40 | 3,516.4 | \$48.84 | 0 | 0.0 | 784.4 | 0.0 | 184.9 | 10 | |
| THURSTON | 5 | 33.4 | \$215.55 | 33.4 | 0.0 | 0.0 | 33.4 | 33.4 | 8 | |
| WAHIAKIUM | 13 | 374.4 | \$158.56 | 87.6 | 286.8 | 0.0 | 273.8 | 374.4 | 40 | |

| Washington | | U.S. Department of Agriculture - Farm Service Agency | | | | | As of: 08-29-03 | | | |
|-----------------------|------------------------|--|------------------|-----------------------|---------------------------|-----------------------|--------------------------|---------------------|--------------------|--|
| Report ID - MEPRTN-R1 | | Conservation Reserve Program - Monthly Contract Report | | | | | Prepared on: 08-29-03 | | | |
| | | Summary for Active Contracts for all Program Years (1986-2004) | | | | | Page: 13 | | | |
| County Name | Total No. of Contracts | Total CRP Acres | Avg. Rental Rate | Continuous CREP Acres | Continuous Non-Crep Acres | Wetland Systems Acres | Margin Pastureland Acres | Tree Practice Acres | Avg. Erosion Index | |
| WALLA WALLA | 539 | 149,966.2 | \$53.06 | 1,501.20 | 2,573.0 | 0.0 | 1,496.9 | 1,728.1 | 10 | |
| WHATCOM | 87 | 1,021.5 | \$347.06 | 1,021.50 | 0.0 | 0.0 | 858.3 | 1,021.5 | 1 | |
| WHITMAN | 1,720 | 138,802.3 | \$74.16 | 0 | 32,203.6 | 456.5 | 754.0 | 1,061.7 | 12 | |
| YAKIMA | 185 | 53,341.3 | \$39.58 | 147.2 | 497.7 | 0.0 | 235.0 | 235.0 | 10 | |
| STATE TOTAL: | 9,878 | 1,286,822 | \$52.14 | 7,080 | 83,609 | 7,906 | 13,880 | 17,953 | 8 | |

<http://www.fsa.usda.gov/crpstorpt/08Approved/r1sumyr/wa.htm>

Appendix E

Upper Middle Mainstem Columbia River Subbasin

Water Quality Parameters Affected by Hydropower Production

Total Dissolved Gas

Total dissolved gas (TDG) supersaturation often occurs during periods of high runoff and spill at hydropower projects and can be harmful to fish. Supersaturation occurs when gases, entrained by water passing over spill gates, are carried to depth by the plunging action of the spill and forced into solution by increased hydrostatic pressure (Perleberg and McDonald 2000). Fish and other aquatic organisms that are exposed to excessive TDG supersaturation can develop gas bubble trauma (GBT), a class of harmful and potentially fatal symptoms. Total dissolved gas supersaturation in the Columbia River was identified in the 1960's and 1970's as a potential detriment to salmon. Those concerns have reappeared as management agencies have reinstated spill as a means of aiding downstream fish passage throughout the system.

The WDOE has set a TDG standard of 110 percent of saturation for all flowing waterways. The WDOE has approved an interim modification to the standard of 110 percent to allow spill for fish passage. The revisions under this modification to state water quality standards allow an average TDG level of 120 percent for the highest 12 hours of a day at the tailrace of the respective dam and allow an average of 115 percent for the highest 12 hours of the day at the forebay of the next downstream dam. The modification to state water quality standards also incorporates a maximum one-hour average TDG reading of 125 percent in the tailrace. These standards do not apply during periods when the river flow exceeds the seven-day, 10-year-frequency flood (7Q10-the level of a flood release that could be expected to occur for a period of seven days on the average of once in ten years). Total dissolved gas at the UMM hydro projects is monitored in both the forebay and tailrace of the projects. The projects typically remain in compliance with the WDOE standards, but on occasion, TDG levels exceed the maximum allowed. This exceedance usually occurs during periods of high run-off or when the water coming into a project is nearing, or is out of compliance with WDOE standards.

Water Temperature

The effect of hydropower projects on Columbia River water temperature has been to delay the time when thermal maximums are reached and when cooling begins in late summer (BPA et al. 1994). The thermal regime of the UMM is largely influenced by releases from Grand Coulee Dam, which is the main upstream deepwater storage project. The UMM hydroelectric projects are run-of-river facilities with very limited capability for storage and flow regulation. In general, the low retention times of the reservoirs at these facilities limit the potential warming that can occur.

Dissolved Oxygen

Dissolved oxygen (DO) levels in the subbasin do not typically decline below the minimum Environmental Protection Agency (EPA) standard for DO in Class A waters of 8.0mg/l.

Turbidity and Suspended Sediments

Turbidity and suspended sediments in the UMM are relatively low (BPA et al. 1994). The hydroelectric projects and their associated reservoirs slow the river flow and allow sediment to settle out. Turbidity and suspended sediments are commonly higher in the tributaries than in the Columbia River of the Columbia River (BPA et al. 1994).

Nutrients

Water quality stations throughout the Columbia River typically show ammonia concentrations that are below the EPA chronic freshwater standard. Mean annual phosphate concentrations often exceed levels that could stimulate algal blooms. Highest phosphate levels occur at the start of spring runoff, and in the late fall at the end of the low-flow season. High levels are also encountered in winter when biological uptake is lowest (BPA et al. 1994).

High levels of nitrates and phosphates have also been observed in the upper reaches of the Douglas Creek watershed, the main tributary to the Moses Coulee (Isaacson 1989), though water samples from lower reaches show higher water quality. Bartu and Andonaegui (2001) suggested that the higher flows in the lower reaches of Douglas Creek might be acting to dilute the levels of nitrates and phosphates.

Appendix F
Upper Middle Mainstem Columbia River Subbasin

**CONSIDERATIONS FOR MONITORING
IN SUBBASIN PLANS**

FROM THE

**PACIFIC NORTHWEST AQUATIC MONITORING
PARTNERSHIP**

May 4, 2004

Table of Contents

| | |
|---|----|
| Introduction | 1 |
| Overview of the Pacific Northwest Aquatic Monitoring Partnership | 2 |
| Nexus with Subbasin Planning | 3 |
| Limitations of This Guidance | 5 |
| Assumptions Regarding Development of Monitoring Elements of Subbasin Plans | 6 |
| General Considerations for Creating Monitoring and Evaluation | |
| Elements of Subbasin Plans | 7 |
| A Strategic Monitoring Framework for Subbasin Planning | 7 |
| Principles for Coordinated Monitoring | 8 |
| Summary of General Considerations | 13 |
| Specific Considerations Regarding the ISRP Review Checklist | 15 |
| Monitoring Objectives | 15 |
| Monitoring Indicators | 16 |
| Data and Information Archive | 17 |
| Coordination and Implementation | 18 |
| RME Logic Path (Evaluation and Adaptive Management) | 19 |
| General Considerations for Creating Monitoring and Evaluation | |
| Elements of Subbasin Plans | 21 |
| Relationship of Subbasin Plans to Existing Monitoring Efforts | 21 |
| Suggested Table of Contents | 24 |
| Program Setup | 24 |
| Detailed Technical Considerations Supporting the Table of Contents | 25 |
| Basic Statistical Considerations | 25 |
| Sampling Design Considerations | 25 |
| Spatial Scale | 26 |
| Classification | 27 |
| Indicators | 28 |
| Measuring Protocols | 29 |
| Status/Trend Monitoring | 31 |
| Appendix A- Participants in PNAMP | 33 |
| Appendix B - Examples of Key Monitoring Questions | 34 |
| Literature Cited | 37 |
| Relevant References | 40 |

Acknowledgements

This document resulted from the collective effort and contributions of many individuals involved in the Pacific Northwest Aquatic Monitoring Partnership. The following people (in alphabetical order) provided text and/or commented on draft materials: Jen Bayer and Dave Busch (U.S. Geological Survey), Bruce Crawford (Washington Salmon Recovery Funding Board and Interagency Committee for Outdoor Recreation), Scott Downie (California Fish and Game), Jim Geiselman (Bonneville Power Administration), Steve Katz, John Stein and Stewart Toshach (NOAA Fisheries Northwest Fisheries Science Center), Steve Lanigan (Forest Service), Steve Leider (Washington Governor's Salmon Recovery Office), Kelly Moore (Oregon Watershed Enhancement Board), Michael Newsom (Bureau of Reclamation), Dave Powers (Environmental Protection Agency), Phil Roger (Columbia River Inter-Tribal Fish Commission), Bruce Schmidt (Pacific States Marine Fisheries Commission), Steve Waste (Northwest Power and Conservation Council), Keith Wolf (Colville Tribes), and Frank Young (Columbia Basin Fish and Wildlife Authority).

Introduction

At the request of the Northwest Power and Conservation Council (Council), the Pacific Northwest Aquatic Monitoring Partnership (PNAMP) has developed this guidance to help subbasin planners design the monitoring elements of the subbasin plans being developed under the Council's Fish and Wildlife Program. It provides general and some specific considerations to the Council and subbasin planners on how their monitoring can fit within the broad range of monitoring activities in the Pacific Northwest. It also provides an explanation of general technical considerations for implementing the various types of monitoring and related topics.

PNAMP offers this initial guidance for monitoring efforts at the subbasin level as a step to encourage the coordination of local, tribal, state and federal programs. Subbasin planners can decide to whether or not, and to what degree, they may elect to use this guidance. PNAMP understands that this guidance is being offered very late in the planning process and therefore does not intend that it add new requirements, but rather that it provide near-term guidance to those still formulating or modifying the monitoring elements of their subbasin plans. This guidance will be less useful to those subbasin planners who are well along in the development of the monitoring elements of their plans, but should nonetheless provide information for those who may modify their plans at a later time.

Overview of the Pacific Northwest Aquatic Monitoring Partnership

Monitoring efforts have typically evolved in response to different organizational mandates and management questions. Despite inherent differences much overlap exists across broad geographical areas, and there are issues and questions shared in common. Collecting monitoring data in a fashion that can be "rolled-up" to larger scales is essential for information gathered at the scale of watersheds or subbasins to support evaluations at larger geographic scales, such as province or Evolutionarily Significant Unit (ESU). This necessitates a higher level of coordination and creates a new set of challenges at all levels of involvement. Toward that end, the PNAMP drafted a coordination plan for monitoring in the Pacific Northwest, *"Recommendations for Coordinating State, Federal, and Tribal Watershed and Salmon Monitoring Programs in the Pacific Northwest"* (PNAMP 2004).

The purpose of PNAMP "is to coordinate monitoring of important scientific information at the appropriate scales needed to inform public policy and resource management decisions" (PNAMP 2004). Members of PNAMP include state, federal, and tribal representatives with a common interest in regionally coordinating various aspects of watershed condition monitoring, fish population monitoring, action effectiveness monitoring, and data management (see Appendix A - Participants in PNAMP). The current focus of PNAMP is on watershed condition and anadromous fish. PNAMP has not made a decision at this time on whether to coordinate monitoring of resident fish and wildlife in the future. Consequently, the scope of this document is limited to monitoring of watershed condition and anadromous fish, and it does not address monitoring of resident fish and wildlife. Subbasin planners can consider the guidance developed by

Council for monitoring these species, as provided in the *Technical Guidelines for Subbasin Planners* and other documents.

Nexus with Subbasin Planning

In January and February of 2004, PNAMP provided briefings to the Council and other regional state, tribal and federal executive level groups on its draft coordination plan. At their briefing to the Council's Regional Coordination Group (RCG) for subbasin planning, PNAMP was asked to provide what guidance it could in the limited time available to assist subbasin planners in developing the monitoring elements of their subbasin plans. In response to that request, PNAMP is herein providing the Council and subbasin planners with guidance and considerations for monitoring. This guidance is advisory in nature, as PNAMP has no inherent authority. PNAMP is an ad hoc collaborative group currently operating without funding or charter that is motivated by the need for technical coordination between its members and across various programs. Despite these limitations, the group elected to provide guidance because several members of PNAMP are involved with subbasin planning, and because the Columbia River Basin constitutes a sizable portion of the geographic scope of PNAMP, from Canada to Northern California. In sum, it is not the intention of PNAMP to dictate a particular direction to subbasin planners, but rather to share the current thinking of the group on many topics relevant to the development of monitoring elements of subbasin plans.

In 2000 the Council initiated subbasin planning to help local entities work with resource experts and managers to develop their own restoration plans. Subbasin planning incorporates a bottom-up approach, with input from a wide range of stakeholders and professionals who are most familiar with the logistical needs in their areas. The Council has stipulated that subbasin plans include a monitoring element. (Monitoring is also required in salmonid recovery plans.) The Council requirements for the monitoring components of subbasin plans were first provided two years ago in the *Technical Guidance for Subbasin Planners* (NPPC 2001). Although subbasin planning remains a bottom-up initiative, several developments within the field of monitoring and data management over the last two years have shifted the Council's perspective on the efficacy of the bottom-up approach for monitoring.

Programmatic or Regional Approach: The need for more extensive, programmatic level habitat and fish population performance tracking and action effectiveness research have emerged as critical elements of survival and recovery plans for salmonids listed under the Endangered Species Act (ESA). Consequently, monitoring questions have been identified in the Federal Salmon Recovery Strategy and the Implementation Plan of the Action Agencies addressing the NOAA-Fisheries Biological Opinion (Biological Opinion) on the Federal Columbia River Power System (FCRPS). (Note: the Action Agencies are Bonneville Power Administration, the Army Corps of Engineers, and the Bureau of Reclamation.) The monitoring questions now being asked are best answered at large-scale landscape and ecosystem levels. The Federal Research, Monitoring and Evaluation Plan for the FCRPS Biological Opinion and the detailed Upper Columbia Monitoring Strategy document the need for this approach. Monitoring and evaluation is

also required under the Pacific Coastal Salmon Recovery Fund. Furthermore, scientific reviews by the Independent Scientific Advisory Board and the Independent Scientific Review Panel have repeatedly called for a regionally coordinated approach to monitoring. Although the Council has reaffirmed the bottom-up approach in regard to other aspects of subbasin planning, the RCG has acknowledged the importance of developing a regional approach to monitoring that will support planning and the setting of restoration priorities across different geographic scales. This is a long-term need of the Council's Fish and Wildlife Program and an immediate need for ESA planners across the Pacific Northwest. One of the tasks of PNAMP is to identify the common metrics and designs necessary to address questions at and across these different scales.

Subbasin or Project Approach: PNAMP intends that this initial guidance constitute a first step in an on-going effort to support local programs in the Pacific Northwest as a means to grow a coordinated regional monitoring program over time. A majority of monitoring work is still occurring at the project scale, for example, in support of individual habitat projects. Yet, comprehensive monitoring strategies consistent with the federal initiatives are now being implemented at the state level in Oregon and Washington. Pilot projects are currently being implemented or planned in the Wenatchee, John Day, and Upper Salmon rivers to collect data and to test and develop more precise protocols and provide increasingly explicit guidance based on field-tested approaches at the subbasin level. (These pilot projects demonstrate how the top-down approach can work to create monitoring projects that have systemwide applications.)

For these reasons, it is clear that both bottom-up and top-down approaches are necessary to develop effective and efficient monitoring plans across the Pacific Northwest. PNAMP sits squarely in between a network of executives who administer resource management programs (top level) and PNAMP members and their constituent groups who implement restoration projects in support of these programs (bottom level). Thus, PNAMP is in the middle, coordinating the most effective system design and application of individual or local projects, such as the pilot studies and NMFS's trend monitoring project.

Collaborative Approach: The progress that PNAMP has made over the last several years is in large measure a result of its collaborative mode of operation. PNAMP is working to coordinate existing monitoring programs and to address issues that challenge practitioners of monitoring irrespective of their geographical location or jurisdictional mandate. PNAMP is not a planning forum or a program, but rather a technical work group whose primary incentive for coordination is the efficiencies to be gained through working collaboratively.

PNAMP, with its mission to improve coordination of monitoring across multiple regional monitoring and evaluation programs, recognizes the importance and challenges of coordinating across the many subbasin monitoring and evaluation plans. If these plans are not coordinated it will be very difficult to add up the results across multiple plans and make conclusions at broader scales, for example at the population level. PNAMP recognizes that while helping monitoring programs in the Pacific Northwest strive to

achieve a greater degree of coordination there will be difficulty in making changes in ongoing monitoring programs. Yet subbasin planning presents PNAMP with an opportunity not unlike that of the Pacific Coastal Salmon Recovery Fund, in which a subset of members have a specific goal, the achievement of which is beneficial to the parent group.

It is important that PNAMP continue to develop technical tools and methodologies that are useful at different scales and for multiple efforts across constituent groups. PNAMP will endeavor to develop additional products for use in the Pacific Northwest that subbasin efforts can use for 2005 and later field seasons. PNAMP members have previously called for workshops on various topics of interest to its members across the Pacific Northwest Region. If these workshops are held (sometime after the subbasin planning submission deadline), they would benefit from the participation of subbasin planners.

PNAMP Coordination Plan: PNAMP intends to complete work on its coordination plan, by fashioning it into a forward-looking, Strategic Monitoring Framework. The exercise of completing the PNAMP plan will provide Tribal and State representatives to PNAMP a better vehicle for coordinating with subbasin planners into the future than this guidance can provide, since it is a response to a Council request for immediate assistance. PNAMP has long provided a forum for coordination amongst its current members, who number over thirty entities representing a broad array of entities and geographic areas. In light of the number of watersheds in the Columbia River Basin (62) and the even larger number between the Canadian border and Northern California, PNAMP members who represent state monitoring programs along with subbasin coordinators, will provide the initial points of contact for subbasin planners and PNAMP. During the implementation of subbasin plans in the Columbia River Basin, PNAMP can be viewed as a source of technical expertise on monitoring in the Pacific Northwest.

Limitations of This Guidance

The PNAMP guidance is divided into sections explaining general and specific considerations. The latter section outlines current PNAMP thinking and experience in regard to relevant technical issues. Please note that some of these considerations may change over time as this coordination effort develops further. Because the Council's *Technical Guidance for Subbasin Planning* (NPCC 2001) states, "the monitoring plan should not include project specific monitoring," this guidance does not address considerations for monitoring at the project scale.

PNAMP accepted the task of helping subbasin planners because of the significant opportunity it afforded to improve coordination of regional monitoring efforts. Despite the very tight deadlines with which subbasin planners are confronted, PNAMP has tried to provide the best guidance possible in the time available. However, PNAMP fully recognizes that the guidance has limitations. The guidance is not sufficiently detailed to represent a complete step-by-step "how-to-guide" or tutorial for monitoring, nor is it based (as would be desired) on a survey of all subbasin planning needs. However,

PNAMP feels it represents a “checklist” of critical elements and other considerations for use in developing subbasin monitoring efforts, and it offers direction for access to example protocols.

This guidance is not intended to supplant the efforts of subbasin planers who are well along in the development of the monitoring elements of their plans. Rather, PNAMP hopes to provide guidance for these efforts and other similar efforts into the future, while providing near-term guidance to those still formulating the monitoring elements of their subbasin plan.

Assumptions Regarding Development of Monitoring Elements of Subbasin Plans

1. Monitoring and evaluation coordination and implementation will be an ongoing activity at the reach, subbasin, and regional levels. PNAMP assumes these iterative, concurrent processes at different scales will be coordinated to optimize when and where implementation occurs to increase learning from broader scale monitoring both within and across subbasins. It is important to note that PNAMP provides a coordination function; PNAMP itself will not implement monitoring.
2. Monitoring that is proposed will be more effective if it fits within a broader programmatic network of status monitoring programs and intensively monitored watersheds. PNAMP assumes subbasin efforts will be able to rely on the broader monitoring framework and programmatic activities to meet some of their needs.
3. PNAMP assumes local, bottom-up approaches developed within subbasins will have higher likelihood for successful funding and meaningful results if they reflect the approaches being developed within the comprehensive state, tribal initiatives, and federal pilot projects (Wenatchee, John Day, and Upper Salmon), and the top-down framework and considerations being developed by PNAMP.
4. PNAMP assumes monitoring elements of subbasin plans that diverge from PNAMP guidance will be explained and framed as pilot approaches to address uncertainties in monitoring strategies or protocols.
5. Additional coordination issues pertaining to larger spatial scales will be identified through PNAMP efforts.

General Considerations for Creating Monitoring and Evaluation Elements of Subbasin Plans

A Strategic Monitoring Framework for Subbasin Planning

The considerations in this section will help the Council and subbasin planners determine the appropriate scales of monitoring and evaluation needed to meet the vision, goals and objectives of subbasin plans. It provides an approach that can be voluntarily used as a foundation for a more detailed, regionally compatible monitoring and evaluation plan.

The implementation and adaptive management of subbasin plans will be difficult absent a well-developed and consistent monitoring framework for the region. The draft PNAMP monitoring coordination plan is intended to develop regional-level guidance for use by the various programs of the members. PNAMP recommends that the implementation of monitoring program elements identified through subbasin planning (bottom-up) be consistent, to the extent practical, with the draft PNAMP plan for coordinating monitoring across the Pacific Northwest (top-down) and recognizes the necessity of both. Conceptually, PNAMP's support for a hierarchical approach to monitoring is linked to guidance provided by the FCRPS Biological Opinion RME Plan and monitoring strategies developed by Oregon and Washington (Table 1). In general, PNAMP sees a role for monitoring within the subbasins with respect to documenting implementation of restoration actions. Subbasin and ESU scale status and trend monitoring are likely to be the responsibility of agency programs that will also need coordination. Evaluating the effectiveness of federal, tribal, and state programs will require participation and cooperation of all those involved with responsibility for evaluation of the plans (Table 2).

PNAMP is working to coordinate current regional monitoring programs that overlap one another at various spatial and temporal scales. Those programs include:

- Aquatic and Riparian Effectiveness Monitoring Program for the Northwest Forest Plan (AREMP);
- Pacfish/Infish Biological Opinion for the interior Columbia Basin (PIBO) Program;
- Interior Columbia Basin Ecosystem Management Program (ICBEMP);
- Columbia River Research, Monitoring, and Evaluation (RME) Program required by ESA Columbia River Biological Opinions and the Columbia River Federal Salmon Recovery Strategy MOU;
- EPA's Environmental Monitoring and Assessment Program;
- NOAA's Pacific Coastal Salmon Recovery Fund Program;
- Monitoring programs associated with salmon recovery and watershed restoration in Oregon, Washington, California, and Idaho;
- National Park Service Monitoring Program;
- Collective and individual tribal monitoring programs; and,
- Co-manager harvest and hatchery monitoring programs.

PNAMP expects to develop further information that should greatly aid monitoring coordination within the Columbia River Basin and across the entire Pacific Northwest.

Over the next year PNAMP will draft a Strategic Monitoring Framework that will identify:

1. A watershed condition and fish population status-monitoring network;
2. A network of Intensively Monitored Watersheds (IMWs) to monitor the effectiveness of different categories of actions on fish at watershed scales; and,
3. Linkages among an identified suite of local, reach specific, action effectiveness studies.

The Strategic Monitoring Framework will identify resources across the cooperating agencies that can help implementers of the subbasin plans to appropriately scale, design and fund their programs. In regards to watershed condition and fish population status monitoring, it is expected that this expanding network of monitoring programs will also lead to research relevant to a majority of the subbasins, including the identification of local, spatially, or temporally intensified monitoring needs. PNAMP suggests that subbasin plans identify their status monitoring needs as:

1. Relying upon work conducted under an existing monitoring program wherever possible;
2. A component of, or cooperator in, an existing monitoring program;
3. A needed addition under an existing or planned program; or
4. An independent, cooperating, contributor to the network of programs.

The federal Action Agencies are implementing three subbasin pilot studies as part of the requirements of the FCRPS Biological Opinion. The state of Washington is initiating IMW efforts that include work in the lower Columbia River. The Bonneville Environmental Foundation is also sponsoring a ten-year program for three IMWs. PNAMP suggests that subbasin plans indicate whether their subbasin is now designated as a subbasin pilot or an IMW, or whether planners think it may serve as a good candidate for this type of monitoring. Viable candidates for IMWs should have characteristics amenable to experimental design features as well as a reasonable potential for management manipulations involving monitoring at multiple treatment and control sites for different categories of individual or combination of actions across an entire watershed. IMWs depend on reliable and precise sampling of adult spawners and smolt outmigrants.

Principles for Coordinated Monitoring

As described in PNAMP (2004), monitoring involves the deliberate and systematic observation, detection, and recording of conditions, resources, and environmental effects of management and other activities. The clear articulation by policy makers of guiding principles helps partners recognize program elements and objectives they share in common. Although PNAMP's draft coordination plan for monitoring addresses an area of greater geographic scope than the Columbia River Basin, its principles may be useful to subbasin planners as they develop the monitoring element of their plans. PNAMP's

Table 1. Example of PNAMP Strategic Framework for Monitoring and Evaluation currently under development: Overview of spatial and temporal scale for monitoring activities with example monitoring types and indicators. Suggested protocols and funding sources under evaluation by PNAMP are included.

| <u>Spatial Scale</u> | <u>Frequency</u> | <u>Monitoring Type</u> | <u>Key Indicators</u> | <u>Protocols to Consider</u> | <u>Potential Funding Sources</u> |
|---|--|--|---|---|---|
| Region – Wide States Major Basins | Infrequent Depending on Activity | Broad Scale: Remote Sensing, Qualitative Surveys, GIS Analysis | Land Use, Roads, Ownership, Vegetation Fish Presence, Intrinsic Habitat Potential | Established Protocols Enhanced Analysis | Existing Federal, State and local government programs. |
| Subbasin ESU Oregon Plan Report Area | Annual and/or Seasonal: Ongoing Duration | Status and Trend Spatially explicit, Rigorous, statistical sampling designs and protocols | Population Abundance, Distribution, Diversity Watershed Condition Riparian & Channel Habitat, Water Quality & Biotic Indicators | EMAP Based Sample Site Selection: Site specific activities (Upper Columbia Monitoring Strategy, AREMP, PIBO, Habitat, Water Qual., Fish Populations, etc) | State F&W or WQ Programs, BPA Fish Program, Action Agencies, NOAA, PCSRF, AREMP, other. |
| Watersheds 5 th -6 th Field (USGS HUC) WA WIRAS | Seasonal and Continuous: Long Term Duration (10-40+ yrs) | <u>Intensively Monitored Watersheds</u> Limiting Factors, BMP Evaluation & Compliance, Effectiveness | Landscape Assessment Watershed Condition and Processes, Salmomid Freshwater Survival & Productivity Management Actions | Upper Columbia Monitoring Strategy, CLAMS, AREMP, Current WA and OR IMW's. Paired-watersheds and/or sample-based watersheds | BPA, NOAA, PCSRF, States & Tribes, Landowner University Co-ops, USFS, BLM, others |
| Stream Reaches | Annual and Seasonal: Med. Duration (5-10 yrs) | <u>Project Effectiveness</u> Desired physical and biotic responses. | Channel and Riparian Habitat Response Fish Use / Productivity Water Quality | Upper Columbia Monitoring Strategy, OPSW Water Quality and Riparian Guides, | BPA, OWEB, SRFB, NOAA and other Funding Entities |

| | | | | | |
|----------|-----------------------------------|--|--|--|---|
| Sites | Seasonal: Short Term (1-5 yrs) | <u>Validation</u> Testing Restoration Methods | Expected vs. Response Conditions. What works, why, and where? | WA-SRFB Protocols, Ongoing PNAMP Process, etc. Various: See WA SRFB Draft Protocols, Upper Columbia Monitoring Strategy | BPA, PCSRF, OWEB, SRFB, other Funding Entities |
| Projects | Before/After Project Completion | <u>Implementation</u> | Location, Description of Activity, Target Species, Ecosystem Function or Habitat Condition | Documentation & Reporting via BiOp Implementation Plans, PRISM, OWEB, NOAA | Condition of Restoration Contract Acceptance (BPA, OWEB, SRFB, USFWS, others) |

Table 2. Generalized Description of Sampling Approaches with Comparative Level of Federal, Tribal, State, and Local Participation.

| <u>Sampling Approach</u> | <u>Spatial Scale</u> | <u>Monitoring Type</u> | <u>Who Does the Work?</u> |
|--|---|--|--|
| Comprehensive Low Intensity | Region – Wide State - Wide Major Basins | Broad Scale Remote Sensing and Surveys | NOAA, Contractors, University, State, Fed and Tribal GIS Programs |
| Sample Based Every Subbasin (~10-30% of fish distribution) | Subbasin ESU Oregon Plan Report Area | Status and Trend | Action Agencies State Agencies Tribes |
| Sample Design or Opportunity Ideally 1 or 2 IMW's In Each Subbasin, ESU, or Report Area | Watersheds 4 th , 5 th , 6 th Field USGS HUC WA WIRA's | Intensively Monitored Watersheds | Multi-Agency University Research Lead Entities Land Owner or Manager |
| Sample Based Stratified by Activity & Location (~20% of Projects) | Stream Reaches | Project Effectiveness | Agencies Research Entities Lead Entities |
| Sample Based Stratified by Project Type (~10% of each type) | Sites | Technical Validation | Lead Entities Agencies |
| Every Project | Projects | Implementation | Subbasin Lead Entities Grantees |

principles include several directives for its members that subbasin planners are encouraged to consider. These principles are:

1. Resource Policy and Management: The purpose of monitoring efforts is to provide the most important scientific information needed to inform public policy and resource management decisions.

- Acknowledge each party's mandates, objectives, and management milestones.
- Construct a monitoring program that meets each party's milestones and objectives through coordinating and sharing monitoring resources.
- Develop a monitoring program that is sufficiently robust to meet public policy needs; demonstrate the links between public policy needs and monitoring efforts.
- Develop a monitoring program that demonstrates compliance.
- Commit to resolving scientifically the most important policy and management questions using an adaptive management approach.

2. Efficiency and Effectiveness: Cooperative monitoring will enhance efficiencies and effectiveness of our respective and collective efforts.

- Participate fully in the PNAMP, including the identification of contact(s) for monitoring issues.
- Identify and coordinate goals, objectives, and budgets, and demonstrate resource savings over short and longer time frames.
- Cooperatively adapt programs and budgets to address monitoring gaps.
- State and federal agencies and the tribes commit to long term inter- and intra-agency monitoring programs.
- Encourage staff exchanges and shared training to learn what each other are doing (e.g., new innovations) and ensure consistency across programs.
- Develop common monitoring approaches, including quality control/quality assurance programs; shared evaluation tools; integrated status and trend monitoring efforts; land use, land cover, and riparian vegetation categorization; core data for representative subset of watersheds in all represented states.
- Perform all monitoring activities in a timely manner.

3. Scientifically Based: Environmental monitoring must be scientifically sound.

- Develop an integrated monitoring program (e.g., issues, disciplines, and values).
- Monitoring program is based on shared goals and objectives (e.g., census level, regional status and trends, cause and effect questions, effectiveness of regional efforts, identification of trouble spots).
- Address multiple spatial and temporal scales.
- Develop and use compatible data collection and analysis protocols.
- Recognize inherent diversity and variability and dynamic inter-relationships or resource conditions in monitoring design, analysis and interpretation.
- All environmental data should have a known level of precision.
- All baseline data on ecosystems are known and compiled between agencies.

- 4. Shared Information:** Monitoring data should be accessible to all on a timely basis.
- Make strategic investments in information systems needed to make data useful.
 - Monitoring databases would integrate a number of issues, disciplines and values.
 - Data management systems and protocols provide a linkage for sharing data between agencies.
 - Adopt and use common data sharing protocols.
 - Adopt and use common database/s of core metadata, data, and electronically connected distribution systems.

Summary of General Considerations

1. It is important to first identify the management questions that any monitoring program is intended to address. (Appendix B provides examples of management questions that are the focus of several existing regional monitoring programs.) These broader level questions frame the objectives and scope of a monitoring strategy. Additional, more detailed questions then need to be developed and answered for developing a specific monitoring strategy or program design. (The section on Program Setup can help identify design level questions that need to be addressed in the development of a specific monitoring strategy.)
2. Subbasin plans and their implementation will be significantly strengthened if they incorporate and are consistent with the principles of the draft PNAMP coordination plan.
3. Subbasin plans will be more effective if they establish a method to link with the continuing development of a Strategic Monitoring Framework by PNAMP, and identify and incorporate guidance for local subbasin level monitoring and evaluation that can be incorporated within this framework.
4. Create a process within subbasin plans to incorporate additional guidance from efforts such as the federal Action Agencies' pilot studies, Collaborative Systemwide Monitoring and Evaluation Project (CSMEP), statewide monitoring initiatives, and further PNAMP guidance as it becomes available.
5. Subbasin plans will be more effective if they identify concrete actions and provide specific plans to promote and achieve needed monitoring and evaluation, and are not "plans to do planning."
6. Subbasin plans will be more effective if they identify existing, expanding, or future planned status and trend monitoring programs and action effectiveness research that can be used to partially or completely meet the monitoring and evaluation needs of subbasin plans. (Note: PNAMP has begun to identify the scope of existing monitoring programs.)
7. Subbasin plans will be more effective if they explain how they incorporate existing monitoring guidance from federal, state or tribal programs.

8. PNAMP suggests that local habitat monitoring needs identified in subbasin plans be addressed using procedures and protocols that result in data that can be linked and interpreted at larger spatial scales (e.g., EMAP design, Upper Columbia Monitoring Strategy, and the Washington and Oregon monitoring strategies). This can be achieved by requiring standard monitoring designs and sampling protocols that have been agreed to or that are being compared within the PNAMP process. (Additional technical detail on appropriate fish, action effectiveness, data management and watershed condition sampling protocols will likely begin to be available from PNAMP and others this fall and beyond.)

9. Subbasin plans will be more effective if, to the extent possible, they utilize guidance on specific monitoring standards, protocols and methods as referenced in relevant ongoing efforts or existing documentation.

Specific Considerations Regarding the ISRP Review Checklist

PNAMP understands that the Council's Independent Scientific Advisory Board (ISAB), Independent Scientific Review Panel (ISRP), and Peer Review Groups will be reviewing the subbasin plans. To ensure consistency, the science group reviewers have been provided a checklist, available at:

<http://www.nwcouncil.org/library/isrp/SubbasinPlanReviewGuide.htm>. In this section, PNAMP identifies considerations specific to the monitoring and evaluation elements of the checklist.

Monitoring Objectives (Checklist III.D.2)

PNAMP Consideration 2-1: Adopt a short list of measurable objectives designed to answer subbasin scale questions about the condition of the watersheds and associated imperiled fish. PNAMP recommends that subbasin planners carefully develop the monitoring questions to be answered within the subbasin. After the questions have been developed, they should be prioritized. It is unlikely there will be sufficient funds available to complete all the desired monitoring. Some possible questions include the following examples taken from Washington's Comprehensive Monitoring Strategy (WMOC 2002).

- How are the annual abundance and productivity of salmon by species, ESU, and life stage changing over time within the subbasin?
- What improvements are occurring in restoring the geographic distribution of salmon by ESU, species, and life stage within the subbasin?
- What is the quality of surface waters in the subbasin?
- How are surface water quality conditions changing over time?
- What are the overall impacts of human related activities on freshwater habitat and landscape processes within the subbasin?

Once the monitoring questions have been developed, specific measurable monitoring objectives can be defined to answer the monitoring questions. Following are examples of objectives that tie directly to the monitoring questions given as examples above.

- Measure status and track trends of the numbers of spawning salmon by stock in each subbasin.
- Measure the geographic distribution (identify drainages occupied by salmon) and evaluate trends of salmon in each subbasin. Determine whether their geographic distributions are improving.
- Measure status of identified water quality indicators.
- Measure the trend of identified water quality indicators at stations representing the cumulative effects of human caused impacts and natural conditions.
- Measure status and trends of identified freshwater habitat indicators in the subbasin. Evaluate whether they are improving relative to a desired target or objective

Monitoring Indicators (Checklist III.D.3)

PNAMP Consideration 3-1: Adopt a short list of measurable indicators designed to provide measures of subbasin scale objectives for the condition of the watersheds and associated fish and wildlife. The indicators should be found in commonly accepted protocols where estimates of their variance and coefficient of variation have been obtained, and there is confidence that the indicator can detect change within a reasonable amount of time. Although the PNAMP has not finalized the broader scale strategy and recommended indicators and associated protocols, the currently recommended indicators are described below under the various types of monitoring.

PNAMP Consideration 3-2: Collection of indicator data to meet the objectives of the monitoring program should be implemented using a structured sampling design. The recommended model for development of probabilistic sampling plans for status and trends is the U.S. Environmental Protection Agency's Environmental Monitoring and Assessment Program (EMAP) strategy proposed by the federal Action Agencies and NOAA Fisheries in their "Draft Research, Monitoring and Evaluation Plan for the NOAA-Fisheries 2000 Federal Columbia River Power System Biological Opinion" (The Research, Monitoring and Evaluation Plan, <http://www.efw.bpa.gov/cgi-bin/FW/welcome.cgi>). PNAMP recommends that subbasin planners cooperate with Columbia Basin-wide attempts to develop common probabilistic (statistical) site selection procedures for population and habitat status monitoring. (Information about design approach of EMAP can be found at: <http://www.epa.gov/nheerl/arm/>).

PNAMP Consideration 3-3: PNAMP recommends that status and trends monitoring at the subbasin scale be part of a larger strategy for monitoring regional status and trends. PNAMP agrees with the ISRP that the EMAP probabilistic sampling plan is most

appropriate for estimating status of habitat and fish and for tracking long-term trends in habitat, water quality and fish distribution. PNAMP recommends:

- Developing a regional aquatic monitoring network covering the states of Washington, Oregon, Idaho, and Northern California using the randomized, spatially balanced, probabilistic design developed by the EMAP (Peck et al. 2001). (PNAMP will help facilitate and coordinate this development.) The monitoring network would be flexible to allow reporting of status and trends at various spatial scales (eco-regions, ESUs, subbasins) and across institutional boundaries (i.e., states, tribes, AREMP, PIBO, Interior Columbia Basin). This will facilitate the integration and sharing of multi-agency data collection and interpretation at the broadest scale, statewide, with subbasins participating to add data points complementary to the broader effort and in cooperation with other federal and state efforts and capable of reporting status and trends at subbasin scales, e.g. OWEB, AREMP, and PIBO (Kershner et al., 2001).
- PNAMP will initiate a regional discussion about selecting monitoring sites across the states of the Pacific Northwest, an area within which the Columbia River Basin's 62 subbasins are included, in an effort to encourage individual subbasins toward a scenario where information will be integrated at coarser scales, such as ESUs.
- PNAMP recognizes that subbasin planners and implementers comprise a new and potentially large group of monitoring practitioners in the Pacific Northwest. PNAMP members involved in subbasin planning and implementation can share their experiences with PNAMP, and PNAMP can in turn develop products for its members in the Pacific Northwest that will be useful to subbasin planners. To initiate this interaction PNAMP recommends that a workshop be convened at the earliest opportunity, at which subbasin planners can learn more about the design, rationale, and mechanics of EMAP, and PNAMP members can learn more about the issues ranging across the Columbia Basin.
- The recently completed Pacific Coastal Salmon Recovery Fund (PCSRF) Data Dictionary provides a set of metrics for reporting data concerning the type and extent of salmon recovery work funded under PCSRF, the budget and the organizations involved. Information about projects funded by PCSRF will be accessible at the link: <http://www.nwr.noaa.gov>. (Look under Regional News Releases/Pacific Coast Salmon Recovery Fund for "PCSRF Performance Metrics/Data Definitions Excel spreadsheet 66k.") These metrics are recommended for use in the subbasin efforts to organize and report project level information regardless of funding source, but are not sufficient for reporting scientific data for monitoring and evaluation purposes.

PNAMP Consideration 3-4: PNAMP recommends subbasin planners inventory restoration projects within their subbasins and determine whether the funding entities have provided for reach scale effectiveness monitoring.

PNAMP Consideration 3-5: Monitoring in support of contract compliance is appropriate for individual actions and will need to conform to the requirements of the respective funding agencies.

Data and Information Archive (Checklist III.D.4)

Adequate access to information related to watershed health and salmon recovery is a critical unmet need. The reporting of recovery success depends on consistent data management standards, which in turn can support composite statistics showing cumulative actions of all federal state, tribal, and local entities. The PNAMP data management goal is to: develop or adopt fish and habitat data collection protocols, sampling protocols and analytical methods and, to ensure that data arising from these protocols can be managed, shared and used. There are many different existing interests/initiatives concerned with improving data collection or management in the Pacific Northwest that represents different constituencies, mandates and obligations. There is no common regional data management system of standards or protocols or network that links these interests and initiatives.

PNAMP recognizes a new effort called the Northwest Environmental Data (NED) network (formerly CBCIS) proposes to work within the region to adopt and maintain standards and protocols for data collection and sharing. The role of NED will be to identify, understand, and document where there are gaps and overlaps in collection protocols across the region, and to coordinate efforts to address those gaps and overlaps by identifying where expert work groups are needed. NED may have a key role in support of subbasin plan implementation and information management.

PNAMP Consideration 4-1: PNAMP recommends that subbasin planners not develop separate data management systems for each subbasin. This guidance should help to meet the standards of existing data management systems and to identify mechanisms so that subbasin planners can more easily access these systems. PNAMP recommends that subbasin planners follow a consistent data management methodology that breaks the tasks into distinct steps:

1. Assessing needs and gathering requirements. Understanding the necessary data products, the people who are involved, and when products are needed.
2. Developing a detailed Data Management Coordination Project Plan following forthcoming guidance from PNAMP. Set out the time frame for deliverables, who will do what and when and cost and cost share.
3. Analyzing the requirements. The requirements need to be described in data management terms.
4. To the degree possible, utilize existing database projects and systems.
5. Designing, developing and testing solutions.
6. Transition and training.
7. Deployment.
8. Maintenance.
9. Independent validation and verification.

It is likely that PNAMP will identify coordination and sharing tasks that will require the development and adoption of standard monitoring protocols for both the collection and management of data. The Upper Columbia Monitoring Strategy (UCMS) (Hillman 2004) provides an example of a protocol for collection of data in the field; that is sampling protocols, required variables, etc. Work under the federal pilot projects provides an example of protocols for the management of data, including data definitions, data organization and storage standards.

Coordination and Implementation (Checklist III.D.5)

PNAMP Consideration 5-1: An important goal of PNAMP is to facilitate coordination among monitoring practitioners across the many state and federal monitoring programs in the Pacific Northwest. PNAMP acknowledges that both the degree and the types of monitoring appropriate to implementing the strategies of a particular subbasin may be unique. Further, there are likely to be diverse and not necessarily compatible opportunities for data sharing among proximal monitoring programs. Therefore, PNAMP recommends that subbasin planners and implementers work with the Council and PNAMP to identify and facilitate opportunities for coordination.

RME Logic Path (Evaluation and Adaptive Management)(Checklist III.D.6)

PNAMP Consideration 6-1: Develop the biological vision, objectives, and strategies for the subbasin to be implemented through the management plan. Refer to the specific vision, objectives and/or strategies throughout the plan that tie the subbasin to the larger geographic area of the Columbia Basin and the specific ESUs of the listed species found within the basin. Tie together the monitoring approach to the programs adopted by the state where the subbasin resides, the federal RME plan for the FCRPS Biological Opinion, or recovery plans. The responsibility for decision-making evaluations and management responses is shared by those working on restoration within a subbasin and those working across subbasins.

PNAMP Consideration 6-2: Pilot efforts are an excellent way to coordinate and concentrate support, and explore avenues that may have widespread implications. PNAMP recommends that such work be informed by prior or on-going efforts outside of the subbasin in question. PNAMP is in the process of identifying a network of intensively monitored watersheds (IMWs) or equivalents across the Pacific Northwest. All subbasins do not necessarily need an intensively monitored watershed. PNAMP recommends the subbasins evaluate current monitoring efforts where validation monitoring is occurring or could occur with minimal extra effort or funding. PNAMP recommends IMWs treat specific target species and specific eco-regions. IMWs or equivalents currently under development or being implemented are included in Table 3.

Table 3. Intensively Monitored Watersheds

| Watershed | Species | Funding Entity/Cooperators |
|------------------------------------|--------------------|---|
| Wenatchee River-Upper Columbia, WA | Chinook, steelhead | BPA, BOR, Upper Columbia Salmon Recovery Region, NOAA Fisheries |

| | | |
|--|-----------------------------|---|
| John Day River, OR | Chinook steelhead | BPA, ODFW, NOAA Fisheries, OWEB |
| Clearwater River , ID | Steelhead | Under discussion |
| Lower Columbia (Germany, Mill, Abernathy Creeks), WA | Chinook coho steelhead chum | SRFB, Lower Columbia Salmon Recovery region, WDFW, WECY |
| Hood Canal, WA | Coho steelhead chum | SRFB, Hood Canal Coordinating Council, WDFW, WECY |

For status monitoring, PNAMP anticipates that much of the local need will be met by the expansion of the higher-level network of coordinated programs and recommends relying on and/or identifying how subbasins can contribute to that network of programs. For action effectiveness monitoring, PNAMP is working to coordinate the strategic placement of IMWs noted above that will address the effectiveness of different actions and a limited set of more local, reach specific studies. PNAMP encourages subbasin planners to identify subbasins and associated rationale for their consideration as possible candidate IMWs.

General Considerations for Creating Monitoring and Evaluation Elements of Subbasin Plans

A disciplined, and well coordinated, monitoring and evaluation program is needed to help confirm our scientific assumptions, resolve key scientific uncertainties, and provide the basis for performance tracking and adaptive management. A coordinated program will maximize efficiencies; avoid duplication, and improve experiments to minimize confounding factors or actions.

Relationship of Subbasin Plans to Existing Monitoring Efforts

The technical guidance provided to subbasin planners was helpful, but did not promote the consistent, coordinated monitoring that is needed for the combination and contrast of data at the Tribal Lands, States, Provinces, and Columbia Basin levels. PNAMP suggest that the monitoring sections of individual subbasin plans would benefit if they identify relationships to programmatic and regional or landscape-scale monitoring programs. Therefore, PNAMP suggests that subbasin planners provide the following information on their relationships to monitoring initiatives within the region.

1. A summary table of ongoing monitoring and evaluation activities at the reach, subbasin and watershed level that reports “who, what and where” attributes are urged at a minimum.
2. A short description of how the subbasin plan monitoring element:
 - a. Assesses whether the goals of the subbasin plan are being met, or not;

- b. Contributes to filling critical data gaps in the assessment;
 - c. Complements project effectiveness monitoring; and,
 - d. Describes how subbasin monitoring and evaluation contain complimentary components for measuring regional (e.g., ESU, province or landscape) scale status and trend for fish and wildlife populations.
3. Provides a brief statement about an implementation and coordination strategy.

PNAMP suggests that the following guidance from the Federal RM&E Plan may be useful for framing monitoring and evaluation goals.

1. Track the status of fish populations and their environment relative to required performance standards,
2. Identify the physical and biological responses to management actions,
3. Resolve critical uncertainties in the methods and data required for the evaluation of future population performance and needed survival improvements.

PNAMP suggest that the following guidance for salmon and steelhead may be useful for framing monitoring and evaluation goals.

1. Maintain and modify ongoing monitoring and evaluation efforts until a more structured and coordinated monitoring and evaluation framework and plans are developed and approved.
2. Expeditiously implement monitoring and evaluation actions that address high priority needs.
3. Collaborate with the NMFS recovery planning and research programs, the Federal Caucus' Basinwide Salmon Recovery Strategy, the NWPPC subbasin planning, and State and Tribal planning efforts to develop a basin wide monitoring and evaluation program and data management system.

PNAMP suggest that the following guidance for resident fish may be useful for framing monitoring and evaluation goals.

1. For species such as Kootenai River white sturgeon: define, monitor, and evaluate flows below impediments to meet natural reproduction objectives specified in the final recovery plan(s).
2. For bull trout, to work with the USFWS resident fish recovery planning efforts to obtain basic population and distribution data needed to develop performance standards and to identify critical monitoring and evaluation needs.

PNAMP suggest that the following guidance for developing an implementation and coordination strategy may be useful for framing monitoring and evaluation goal (example from the Oregon Plan).

1. Assess status and trends of watershed conditions and salmon populations regionally.
2. Monitor habitat, water quality, biotic health, and salmon in select watersheds.

3. Analyze habitat, water quality and population trends at the landscape scale.
4. Document conservation and restoration projects, activities and programs.
5. Evaluate effectiveness of restoration and management efforts locally.
6. Evaluate the combined effectiveness of restoration and conservation efforts in select watersheds.
7. Standardize monitoring, collection, management and analysis efforts.
8. Coordinate and support public-private monitoring partnerships.
9. Integrate information and product data products and reports.

The status and trend-monitoring program (NOAA Pilot Studies proposal) for anadromous salmonids and habitat in the Wenatchee and Grande Ronde River basins will serve three major data collection efforts:

- At the scale of a subbasin, assess on an annual basis the status of adult populations of anadromous salmonids.
- At the scale of a subbasin, assess on an annual basis the population status or productivity of juvenile anadromous salmonids.
- At the scale of a subbasin, assess on an annual basis the status of salmonid habitat.

Data from the status and trend-monitoring program will be used for a variety of resource management purposes. The primary utility of the information will be the annual assessment of status and resulting trend over time for these fishes and their habitat. However, monitoring and evaluation programs will also support restoration action planning and assessment by serving as the baseline information used for action siting, and the baseline against the biological impact of actions could be measured.

Other useful references and links include:

1. The Yakima Klickitat Fisheries Project: <http://www.ykfp.org>
2. The Northeast Oregon Hatchery: <http://www.cbfwa.org/2001/projects/198805301.htm>
3. The Columbia Basin Fish and Wildlife Authority (M&E): <http://www.cbfwa.org/rme.htm>
4. The State of Washington: Outline for Salmon Regional Recovery Plans. http://www.wdfw.wa.gov/recovery/recovery_model.htm Coordinated Management Strategy. <http://www.iac.wa.gov/srfb/monitoring.htm>

(Please see the reference sections of this document for a more comprehensive list of resources and full citations.)

This rest of this section is intended to outline considerations for subbasin programs and technical details, intended to facilitate consistency in format and in scientific rigor across subbasins. PNAMP has used the Upper Columbia Monitoring Strategy, or UCMS, (Hillman et al., 2004) as a template for this section because of its current relevancy.

The indicators and metrics contained in the UCMS are derived from NOAA Fisheries, the Federal Columbia River Research and Monitoring and Evaluation (RME) program and component BPA Pilot Projects; the state of Washington's Coordinated Monitoring Strategy, and the Oregon Plan Monitoring Program. Further, detailed guidance in the UCMS incorporates direction and considerations from programs such as: PIBO, AREMP, EMAP, and the WSRFB. Over 35 private, state, federal and tribal representatives have contributed to the development of the UCMS over the course of 2003 and 2004. Thus, the information contained therein, coupled with the following summarized sections, represents the most detailed guidance for program setup, implementation, design, methods, protocols, standards and indicators for monitoring that exist for a Columbia Basin subbasin at this time. Please note that the UCMS also contains many elements and a level of detail that is consistent with an IMW as described previously. However, the UCMS is more detailed than will be needed for all subbasin plans. The UCMS can be accessed online via the Columbia Basin Fish and Wildlife Authority at www.cbfwa.org under the RME section.

The intent of the material that follows is to offer for consideration by planners a concise overview or checklist of steps for development of monitoring plans that would generate statistically valid results. Although these steps are general, PNAMP recommends that planners address each one in order to develop complete understanding of status/trend and action effectiveness monitoring. Below is a suggested table of contents that organizes information according to the steps needed to setup and implement a monitoring program. Following that is an outline of the technical steps needed to effectively design status/trend and action effectiveness monitoring.

Suggested Table of Contents

1. Statement of Need and Program Outline
2. Summary of Indicators and Program Elements
3. Summary of Monitoring and Evaluation Priorities
4. Program Set Up Statistical Design
5. Sampling Design
 - a. Sample Size
 - b. Measurement Error
6. Fish Population Monitoring Overview
7. Habitat Monitoring Overview
8. Biological Variables
10. Physical/Environmental Variables
11. Spatial Scales
12. Performance Standards
13. Classification
14. Indicators to be used
15. Measuring Protocols to be used
16. Status Trend Monitoring
17. Effectiveness Monitoring
18. Data Management Needs Assessment and Data Management Plan

19. Peer Review and Annual Reporting
20. Adaptive Management
21. References
22. Appendices as needed

Program Setup

In order to setup a monitoring program, it will be important to follow a logical sequence of steps. By proceeding through each step, the planner will better understand the goals of monitoring and its strengths and limitations. These steps will aid the implementation of a valid monitoring program that reduces duplication of sampling efforts, and thus overall costs, but still meets the needs of the different entities. The plan assumes that all entities involved with implementing the plan will cooperate and freely share information. Setup steps are:

1. Identify the populations and/or subpopulations of interest (e.g., spring Chinook, steelhead, bull trout).
2. Identify the geographic boundaries (areas) of the populations or subpopulations of interest.
3. Describe the purpose for selecting these populations or subpopulations (i.e., what are the concerns?).
4. Identify the objectives for monitoring.
5. Select the appropriate monitoring approach (status/trend or effectiveness monitoring or both) for addressing the objectives.
6. Identify and review existing monitoring and research programs in the area of interest.
7. Determine if those programs satisfy the objectives of the proposed program.
8. If monitoring and evaluation data gaps exist, implement the appropriate monitoring approach by following the criteria outlined in 9-13.
9. Classify the landscape and streams in the area of interest.
10. Complete a data management needs assessment. Describe how data collection and management needs will be met and shared among the different entities.
11. Identify an existing database for storing biological and physical/environmental data.
12. Estimate costs of implementing the program.
13. Identify cost-sharing opportunities.

Detailed Technical Considerations Supporting the Table of Contents

Basic Statistical Considerations

This document defines “statistical design” as the logical structure of a monitoring study. It does not necessarily mean that all studies require rigorous statistical analysis. Rather, it implies that all studies, regardless of the objectives, be designed with a logical structure that reduces bias and the likelihood that rival hypotheses are correct. The purpose of this section is two-fold. First, it identifies the minimum requirements of valid statistical designs and second it identifies the appropriate designs for status/trend and effectiveness monitoring. The following discussions draw heavily on the work of Hairston (1989),

Hicks et al. (1999), Krebs (1999), Manly (1992, 2001), and Hillman and Giorgi (2002). (See: Hillman et al. 2004) section 3, pages 9-13.)

Sampling Design Considerations

Once the investigator has selected a valid statistical design, the next step is to select “sampling” sites. *Sampling* is a process of selecting a number of units for a study in such a way that the units represent the larger group from which they were selected. The units selected comprise a *sample* and the larger group is referred to as a *population*.¹ All the possible sampling units available within the area (population) constitute the *sampling frame*.² The purpose of sampling is to gain information about a population. If the sample is well selected, results based on the sample can be generalized to the population. Statistical theory assists in the process of drawing conclusions about the population using information from a sample of units.

Defining the population and the sample units may not always be straightforward because the extent of the population may be unknown, and natural sample units may not exist. For example, a researcher may exclude livestock grazing from sensitive riparian areas in a watershed where grazing impacts are widespread. In this case the management action may affect aquatic habitat conditions well downstream from the area of grazing. Thus, the extent of the area (population) that might be affected by the management action may be unclear, and it may not be obvious which sections of streams to use as sampling units.

When the population and/or sample units cannot be clearly defined, the investigator should subjectively choose the potentially affected area and impose some type of sampling structure. For example, sampling units could be stream habitat types (e.g., pools, riffles, or glides), fixed lengths of stream (e.g., 150-m long stream reaches), or reach lengths that vary according to stream widths (e.g., see Simonson et al. 1994). Before selecting a sampling method, the investigator should define the population, size and number of sample units, and the sampling frame. (See: Hillman et al. 2004) section 4, pages 9-13).

Spatial Scale

Because monitoring will occur at a range of spatial scales, there may be some confusion between the roles of status/trend monitoring and effectiveness monitoring. Generally, one thinks of status/trend monitoring as monitoring that occurs at coarser scales and effectiveness monitoring at finer scales. In reality, both occur across different spatial scales, and the integration of both is needed to develop a valid monitoring program (ISAB 2003; AA/NOAA Fisheries 2003; WSRFB 2003).

¹ This definition makes it clear that a “*population*” is not limited to a group of organisms. In statistics, it is the total set of elements or units that are the target of our curiosity. For example, habitat parameters will be monitored at sites selected from the *population* of all possible stream sites in the watershed.

² The *sampling frame* is a “list” of all the available units or elements from which the sample can be selected. The sampling frame should have the property that every unit or element in the list has some chance of being selected in the sample. A sampling frame does not have to list all units or elements in the population.

The scale at which status/trend and effectiveness monitoring occurs depends on the objectives of the study, the size or distribution of the target population, and the indicators that will be measured. In status/trend monitoring, for example, the objective may be to measure egg-parr survival of spring Chinook salmon in the Wenatchee Basin. Because the Wenatchee Basin consists of one population of spring Chinook (ICBTRT 2003), the entire basin is the spatial scale at which egg-parr survival is monitored. In contrast, if the objective is to assess egg-parr survival of spring Chinook in the Chiwawa Basin (a sub-population of the Wenatchee population), the spatial scale at which monitoring occurs includes only the Chiwawa Basin, a much smaller area than the entire Wenatchee Basin. Thus, status/trend monitoring can occur at various scales depending on the distribution of the population of interest.

In the same way, effectiveness monitoring can occur at different spatial scales. That is, one can assess the effect of a tributary action on a specific Recovery Unit or ESU (which may encompass several populations), a specific population (may include several sub-populations), at the sub-population level (may encompass a watershed within a basin), or at the reach scale. Clearly, the objectives and hence the indicators measured dictate the spatial scale at which effectiveness monitoring is conducted. For example, if the objective is to assess the effects of nutrient enhancement on egg-smolt survival of spring Chinook in the Chiwawa Basin (a sub-population of the Wenatchee spring Chinook population), then the spatial scale covered by the study should include the entire area inhabited by the eggs, fry, parr, and smolts. If, on the other hand, the objective is to assess the effects of a sediment reduction project on egg-fry survival of a local group of spring Chinook (i.e., Chinook within a specific reach of stream), then the study area would only encompass the reach of stream used by spawners of that local group.

In theory there might be no limit to the scale at which effectiveness monitoring can be applied, but in practice there is a limit. This is because as the spatial scale increases, the tendency for multiple treatments (several habitat actions) affecting the same population increases. That is, at the spatial scale representing a Recovery Unit, ESU, or population, there may be many habitat actions within that area. Multiple treatment effects make it very difficult to assess the effects of specific actions on an ESU. Even though it may be impossible to assess specific treatment effects at larger spatial scales, it does not preclude one from conducting effectiveness monitoring at this scale. Indeed, one can assess the combined or cumulative effects of tributary actions on the Recovery Unit, ESU, or population. However, additional effectiveness monitoring may be needed at finer scales to assess the effects of individual actions on the ESU or population. (See: Hillman et al. 2004, section 5, pages 31-33.)

Classification

Both status/trend and effectiveness monitoring require landscape classification. The purpose of classification is to describe the “setting” in which monitoring occurs. This is necessary because biological and physical/environmental indicators may respond differently to tributary actions depending on landscape characteristics. A hierarchical classification system that captures a range of landscape characteristics should adequately

describe the setting in which monitoring occurs. The idea advanced by hierarchical theory is that ecosystem processes and functions operating at different scales form a nested, interdependent system where one level influences other levels. Thus, an understanding of one level in a system is greatly informed by those levels above and below it.

A defensible classification system should include both ultimate and proximate control factors (Naiman et al. 1992). Ultimate controls include factors such as climate, geology, and vegetation that operate over large areas, are stable over long time periods, and act to shape the overall character and attainable conditions within a watershed or basin. Proximate controls are a function of ultimate factors and refer to local conditions of geology, landform, and biotic processes that operate over smaller areas and over shorter time periods. These factors include processes such as discharge, temperature, sediment input, and channel migration. Ultimate and proximate control characteristics help define flow (water and sediment) characteristics, which in turn help shape channel characteristics within broadly predictable ranges (Rosgen, 1996).

The UCMS plan proposes a classification system that incorporates the entire spectrum of processes influencing stream features and recognizes the tiered/nested nature of landscape and aquatic features. This system captures physical/environmental differences spanning from the largest scale (regional setting) down to the channel segment. The Action Agencies/NOAA Fisheries RME plan proposes a similar classification system. By recording these descriptive characteristics, the investigator will be able to assess differential responses of indicator variables to proposed actions within different classes of streams and watersheds. Importantly, the classification work described here fits well with Level 1 monitoring under the ISAB (2003) recommend strategies for restoring tributary habitat. Classification variables and recommend methods for measuring each variable are defined below. (See: Hillman et al. 2004) section 6, pages 33-45.)

Indicators

PNAMP has not yet convened a committee to negotiate a set of key indicators for the region. However, a workgroup which includes some PNAMP members has identified the following as a subset of key indicators: bank-full width, reach length, bank-full depth, sediment, wood, gradient, pools, residual pool depth, bank stability, temperature, invertebrates, shade, riparian characteristics. (Please note that this set of attributes has not been reviewed by PNAMP.)

Theses indicators represent a subset of variables that should be measured. Investigators can measure additional variables depending on their objectives and past activities. For example, reclamation of mining-impact areas may require the monitoring of pollutants, toxicants, or metals. Some management actions may require the measurement of

thalweg³ profile, placement of artificial instream structures, or livestock presence. Adding other needed indicators will supplement the core list.

Indicator variables identified in the UCMS template are consistent with those identified in the Action Agencies/NOAA Fisheries RME Plan and with most of the indicators identified in the WSRFB (2003) monitoring strategy. The Action Agencies/NOAA Fisheries selected indicators based on their review of the literature (e.g., Bjornn and Reiser 1991; Spence et al. 1996; and Gregory and Bisson 1997) and several regional monitoring programs (e.g., PIBO, AREMP, EMAP, WSRFB, and the Oregon Plan). They selected variables that met various purposes including assessment of fish production and survival, identifying limiting factors, assessing effects of various land uses, and evaluating habitat actions. Their criteria for selecting variables were based on the following characteristics:

- Indicators should be sensitive to land-use activities or stresses.
- They should be consistent with other regional monitoring programs.
- They should lend themselves to reliable measurement.
- Physical/environmental indicators would relate quantitatively with fish production.

Measuring Protocol

An important component of all regional monitoring strategies (ISAB, Action Agencies/NOAA Fisheries, and WSRFB) is that the same measurement method be used to measure a given indicator. The reason for this is to allow comparisons of biological and physical/environmental conditions within and among watersheds and basins.⁴ This section identifies methods to be used to measure biological and physical/environmental indicators. The methods identified in this plan are consistent with those described in the Action Agencies/NOAA Fisheries RME Plan and, for the most part, consistent with EMAP and WSRFB protocols.

PNAMP is supporting an initiative to coordinate a side-by-side comparison of protocols and will communicate to subbasin planners which protocols will be included in the test. This comparison, which is proposed to take place in 2005, will be done to identify which protocols are best for determining watershed condition status and trend. It's possible a pilot study in the John Day basin will take place in 2004 if funding and logistical constraints are resolved.

The Action Agencies/NOAA Fisheries monitoring group reviewed several publications, including the work of Johnson et al. (2001) that describe methods for measuring indicators. Not surprisingly, there can be several different methods for measuring the same variable. For example, channel substrate can be described using surface visual

³ "Thalweg" is defined as the path of a stream that follows the deepest part of the channel (Armantrout 1998).

⁴ Bonar and Hubert (2002) and Hayes et al. (2003) review the benefits, challenges, and the need for standardized sampling.

analysis, pebble counts, or substrate core samples (either McNeil core samples or freeze-core samples). These techniques range from the easiest and fastest to the most involved and informative. As a result, one can define two levels of sampling methods. Level 1 (extensive methods) involves fast and easy methods that can be completed at multiple sites, while Level 2 (intensive methods) includes methods that increase accuracy and precision but require more sampling time. The Action Agencies/NOAA Fisheries monitoring group selected primarily Level 2 methods, which minimize sampling error, but maximizes cost.

Before identifying measuring protocols, it is important to define a few terms. These terms are consistent with the Action Agencies/NOAA Fisheries RME Plan.

Reach (effectiveness monitoring) – for effectiveness monitoring, a stream reach is defined as a relatively homogeneous stretch of a stream having similar regional, drainage basin, valley segment, and channel segment characteristics and a repetitious sequence of habitat types. Reaches are identified by using a list of classification (stratification) variables. Reaches may contain one or more sites. The starting point and ending point of reaches will be measured with Global Positioning System (GPS) and recorded as Universal Transverse Mercator (UTM).

Although the level of accuracy expected from GPS reporting of stream locations may not be sufficient for all subbasin monitoring and evaluation purposes, the researchers for the John day and Upper Columbia projects are planning to use it for the subbasin pilot efforts.

Reach (status/trend monitoring) – For status/trend monitoring, this section refers only to a “sampling reach” as defined by the EMAP design and referenced in the UC Strategy document. This is one method to consider using to initially locate a reach, with the “X” point being the place where bankfull width is determined. From this location the extent of the upstream and downstream boundaries (total reach length) are determined according to the protocol used. Data collected in the sampling reach should be linked to the best available hydrography layers to facilitate mapping and use in a GIS. Typically the 1:100,000 scale has been used, but a routed 1:24,000 scale hydrography may soon become available.

Note: Standardized GIS and post processing of spatial data will require a standardized protocol that does not currently exist. In the interim PNAMP recommends the following: 1. all GIS data should be provided with Federal Geographic Data Committee compliant metadata, including information on projection used; 2. data should be linked to a standardized stream each identification system to facilitate mapping and use in GIS; and, 3. use existing 1:100,000 and 1:24,000 hydrography layers where they have been cleaned and routed, and if not, use the best available information.

Site (effectiveness monitoring) – a site is an area of the effectiveness monitoring stream reach that forms the smallest sampling unit with a defined boundary. Site length depends on the width of the stream channel. Sites will be 20 times the average bankfull width with a minimum length of 150 m and a maximum length of 500 m. Site lengths are measured along the thalweg. The upstream and downstream boundaries of the site will be measured with GPS and recorded as UTM. For purposes of re-measurements, these points will also be photographed, marked with permanent markers (e.g., orange plastic survey stakes), and carefully identified on maps and site diagrams. Site lengths and boundaries will be “fixed” the first time they are surveyed and they will not change over time even if future conditions change.

Transect – a transect is a straight line across a stream channel, perpendicular to the flow, along which habitat features such as width, depth, and substrate are measured at pre-determined intervals. Effectiveness monitoring sites and status/trend monitoring reaches will be divided into 11 evenly spaced transects by dividing the site into 10 equidistant intervals with “transect 1” at the downstream end of the site or reach and “transect 11” at the upstream end of the site or reach. The number of transects varies for different attributes.

Habitat Type – Habitat types, or channel geomorphic units, are discrete, relatively homogenous areas of a channel that differ in depth, velocity, and substrate characteristics from adjoining areas. This plan recommends that the investigator identify the habitat type under each transect within a site or reach following the Level II classification system in Hawkins et al. (1993). That is, habitat will be classified as turbulent fast water, non-turbulent fast water, scour pool, or dammed pool (see definitions in Hawkins et al. 1993). By definition, for a habitat unit to be classified, it should be longer than it is wide. Plunge pools, a type of scour pool, are the exception, because they can be shorter than they are wide. (See: Hillman et al. 2004) section 8, pages 59-76)

Status/Trend Monitoring

If the objective of the monitoring program is to assess the current status of populations and/or environmental conditions, or to assess long-term trends in these parameters, then the following steps will help the investigator design a valid status/trend monitoring program.

Problem Statement and Overarching Issues:

1. Identify and describe the problem to be addressed.
2. Identify boundaries of the study area.
3. Describe the goal or purpose of the study.
4. List hypotheses to be tested.

Statistical Design (see Section 3 of UCMS Strategy):

1. Describe the statistical design to be used (e.g., EMAP design).
2. List and describe potential threats to external validity and how these threats will be addressed.
3. If this is a pilot test, explain why it is needed.
4. Describe descriptive and inferential statistics to be used and how precision of statistical estimates will be calculated.

Sampling Design (see Sections 4 & 5 of UCMS Strategy):

1. Describe the statistical population(s) to be sampled.
2. Define and describe sampling units.
3. Identify the number of sampling units that make up the sampling frame.
4. Describe how sampling units will be selected (e.g., random, stratified-random, systematic, etc.).
5. Describe variability or estimated variability of the statistical population(s).
6. Define Type I and II errors to be used in statistical tests (the plan recommends no less than 0.80 power).

Measurements (see Sections 7 & 8 of UCMS Strategy):

1. Identify indicator variables to be measured.
2. Describe methods and instruments to be used to measure indicators.
3. Describe precision of measuring instruments.
4. Describe possible effects of measuring instruments on sampling units (e.g., core sampling for sediment may affect local sediment conditions). If such effects are expected, describe how the study will deal with them.
5. Describe steps to be taken to minimize systematic errors.
6. Describe QA/QC plan, if any.
7. Describe sampling frequency for field measurements.

Results:

1. Explain how the results of this study will yield information relevant to management decisions.

Subbasin planners should include a section regarding how the data from the study (with metadata) will be stored, managed and made available to others. A starting point for some subbasin data collection efforts, could be the data definitions document for the Upper Columbia and John Day pilot projects once it has been reviewed. Proponents for the Upper Columbia and John Day projects are reviewing the final data dictionary on which their data system will be developed. The mechanics of data management in the Upper Columbia and John Day systems are being developed by the respective project teams and need significant additional work.

Appendix A - Participants in PNAMP

Tribal

Columbia River Intertribal Fish Commission
Confederated Colville Tribes
Confederated Tribes of the Umatilla Indian Reservation
Northwest Indian Fisheries Commission

State Agencies

California Department of Fish and Game
Idaho Department of Fish and Game
Oregon Department of Environmental Quality
Oregon Department of Forestry
Oregon Department of Fish and Wildlife
Oregon Watershed Enhancement Board
Washington Interagency Committee for Outdoor Recreation
Washington Department of Ecology
Washington Department of Fish and Wildlife
Washington Governor's Salmon Recovery Office

Federal Agencies

Bonneville Power Administration
National Oceanic and Atmospheric Administration
National Park Service
U.S. Army Corps of Engineers
U.S. Bureau of Land Management
U.S. Bureau of Reclamation
U.S. Geological Survey
U.S. Environmental Protection Agency
U.S. Fish and Wildlife Service
U.S. Forest Service
U.S. Park Service

Regional

Columbia Basin Fish and Wildlife Authority
Northwest Power and Conservation Council
Pacific States Marine Fish Commission - StreamNet

Private Sector

BioAnalysts
Bonneville Environmental Foundation
Chelan County PUD
Keith Wolf Associates
Humboldt State University
Paulsen Environmental Research
TetraTech

Appendix B - Examples of Key Monitoring Questions

This section provides selected examples of management level questions that are being addressed under the Washington Comprehensive Monitoring Strategy; the Oregon Plan; the Draft Research, Monitoring and Evaluation Plan for the NOAA-Fisheries 2000 FCRPS Biological Opinion; and the Okanogan Baseline Program.

Washington Comprehensive Monitoring Strategy

1. How are the annual abundance and productivity of salmon by species, ESU, and life stage changing over time?
2. What improvements are occurring in restoring the geographic distribution of salmon by ESU, species, and life stage to their historic range?
3. Are the unique life history characteristics of salmon within a Salmon Recovery Region changing over time because of human activities?
4. What are the trends in the climate of the Pacific Northwest that will allow the State to anticipate and account for such conditions in initiating and monitoring management actions for watershed health and salmon recovery. What trends in climate may mask or expose the status of freshwater habitat and its role in salmon recovery?
5. What are the trends in effects of hatchery production on the survival and productivity of wild salmon populations within each ESU?
6. How are surface water quality conditions changing over time?
7. How effective are clean water programs at meeting water quality criteria?
8. What are the trends in water quantity and flow characteristics?
9. What are the status and trends in habitat-forming landscape processes in riverine tidal, estuarine, and nearshore ecosystems as they relate to watershed health and salmon recovery?
10. Are habitat improvement projects effective?

Oregon Plan for Salmon and Watersheds Monitoring Framework

1. What is the condition of salmon populations at the ESU, Subbasin and watershed scale?
2. What is the status and what are the trends in aquatic habitats, water quality, and stream flow?
3. What are the critical factors that limit watershed function and salmon

productivity?

4. What constitutes detectable and meaningful change in habitat condition and populations?
5. What changes are occurring in watersheds that improve stream habitat quality?
6. What are the management practices and programs that enhance or restore watershed functions and salmon populations?
7. What habitat changes and biotic responses result from these projects, practices, and programs?

**Draft Research, Monitoring and Evaluation Plan for the NOAA-Fisheries
2000 Federal Columbia River Power System Biological Opinion**

1. What are the abundances, productivity, and distributions of Columbia River Basin (CRB) fish populations relative to performance standards or objectives?
2. What are the biological, chemical, and physical status of CRB fish habitat relative to performance standards or objectives?
3. What are the relationships between fish populations and freshwater and estuary/ocean habitat conditions that determine population-limiting factors?
4. What is the effect of a specific mitigation or management action on the habitat and/or population performance of CRB fish?
5. What is the combined effect of multiple watershed level mitigation or management actions on the habitat and/or population performance of CRB fish?
6. Are Federal and state mitigation actions achieving the necessary survival changes identified in the All H Federal Caucus Program and the FCRPS BO for each ESU?

Okanogan Baseline Program - The Colville Tribes (EMAP design):

1. What are the current habitat conditions and abundance, distribution, life-stage survival, and age-composition of anadromous fish in the Upper Columbia Basin (status monitoring)?
2. How do these factors change over time (trend monitoring)?
3. What effects do tributary habitat actions have on fish populations and habitat conditions (effectiveness monitoring)?

4. What effects do fishery management actions have on fish populations (effectiveness monitoring)?
5. Is there is a statistically significant difference in the abundance, survival, and timing and life history characteristics of summer/fall, spring Chinook, sockeye, and steelhead (7-20+ year time frame)?
6. Is there is a statistically significant difference in selected physical habitat parameters and characteristics for summer/fall, spring Chinook, sockeye, and steelhead in the Okanogan basin resulting from the cumulative benefits of habitat actions (7-20+ year time frame)?
7. What is the in-basin and out-of-basin harvest and stock-specific harvest of hatchery and wild anadromous salmonids within the Okanogan subbasin (ongoing)?
8. How effective are selective fishing gears and sites for possible future use for selective Tribal subsistence fisheries?

Literature Cited

- Action Agencies (Bonneville Power Administration, U.S. Bureau of Reclamation, and U.S. Army Corps of Engineers) and NOAA Fisheries. 2003. Draft research, monitoring and evaluation plan for the NOAA-Fisheries 2000 Federal Columbia River Power System Biological Opinion. Bonneville Power Administration, U.S. Bureau of Reclamation, U.S. Army Corps of Engineers, and NOAA Fisheries, Portland, OR. Web link: <http://www.nwr.noaa.gov/1hydrop/hydroweb/fedrec.htm>
- Bonar, S. A. and W. A. Hubert. 2002. Standard sampling of inland fish: benefits, challenges, and a call for action. *Fisheries* 27:10-16.
- Bjornn T. C. and D. W. Reiser. 1991. Habitat requirements of salmonids in streams. *American Fisheries Society Special Publication* 19:83-138.
- Good, T. P., T. K. Harms, and M. H. Ruckelshaus. 2003. Misuse of checklist assessments in endangered species recovery efforts. *Conservation Ecology* 7(2): 12. [online] URL: <http://www.consecol.org/vol7/iss2/art12>
- Gregory, S. V. and P. A. Bisson. 1997. Degradation and loss of anadromous salmonid habitat in the Pacific Northwest. Pages 277-314 in: D. J. Stouder, P. A. Bisson, and R. J. Naiman, editors. *Pacific salmon and their ecosystems, status and future options*. Chapman and Hall, New York, NY.
- Hairston, N. G. 1989. *Ecological experiments: purpose, design, and execution*. Cambridge University Press, New York, NY.
- Hayes, D. and 14 others. 2003. Developing a standardized sampling program: the Michigan experience. *Fisheries* 28:18-25.
- Hawkins, C. P., and ten others. 1993. Hierarchical approach to classifying stream habitat features. *Fisheries* 18:3-12.
- Hawkins, C., J. Ostermiller, M. Vinson, and J. Stevenson. 2001. Steam algae, invertebrate, and environmental sampling associated with biological water quality assessments: field protocols. Utah State University, Logan, UT. Web link: <http://www.usu.edu/buglab/monitor/USUproto.pdf>
- Hicks, L. L., J. Light, G. Watson, B. Sugden, T. W. Hillman, and D. Berg. 1999. Adaptive management: concepts and applications to Plum Creek's Native Fish Habitat Conservation Plan. Native Fish Habitat Conservation Plan Technical Report No. 13, Plum Creek Timber Company, Seattle, WA.
- Hillman, T. W. and A. E. Giorgi. 2002. Monitoring protocols: effectiveness monitoring of physical/environmental indicators in tributary habitats. BioAnalysts, Inc. Report to Bonneville Power Administration, Portland, Or. Web link: <http://www.efw.bpa.gov/cgi-bin/FW/welcome.cgi?ViewMode=External>
- Hillman, T. W. 2004. Monitoring Strategy for the Upper Columbia Basin. BioAnalysts, Inc. Eagle, Idaho. Prepared for: Upper Columbia Regional Technical Team; Upper Columbia Salmon Recovery Board Wenatchee, Washington.
- ICBTRT (Interior Columbia Basin Technical Recovery Team). 2003. Independent populations of chinook, steelhead, and sockeye for listed evolutionarily significant units within the interior Columbia River domain. Working draft. NOAA Fisheries Northwest Fisheries Science Center, Seattle, WA.

- ISAB (Independent Scientific Advisory Board). 2003. A review of strategies for recovering tributary habitat. Independent Scientific Advisory Board for the Northwest Power Planning Council, ISAB 2003-2, Portland, OR. Web link: <http://www.nwppc.org/library/isab/Default.htm>
- Johnson, D. H., and nine others. 2001. Inventory and monitoring of salmon habitat in the Pacific Northwest—directory and synthesis of protocols for management/research and volunteers in Washington, Oregon, Montana, Idaho, and British Columbia. Review draft. Washington Department of Fish and Wildlife, Olympia, WA. Web link: <http://www.wa.gov/wdfw/hab/sshiap/dataptcl.htm>
- Kershner, J. L., E. Cowley, R. Henderson, K. Kratz, D. Martin, C. Quimby, K. Stein, D. Turner, L. Ulmer, M. Vinson, and D. Young. 2001. Effectiveness monitoring of aquatic and riparian resources in the area of PACFISH/INFISH and the Biological Opinions for Bull trout, salmon, and steelhead. Draft plan. USDA Forest Service/USDI Bureau of Land Management. Logan, UT. 50p.
- Krebs, C. J. 1999. Ecological methodology. Second edition. Benjamin/Cummings, Menlo Park, CA.
- Malone, K. Mobrand Biometrics, personal communication.
- Manly, B. F. J. 2001. Statistics for environmental science and management. Chapman and Hall, New York, NY.
- Naiman, R. J., D. G. Lonzarich, T. J. Beechie, and S. C. Ralph. 1992. General principles of classification and the assessment of conservation potential in rivers. Pages 93-123 in: P. J. Boon, P. Calow, and G. E. Petts, editors. River conservation and management. John Wiley and Sons, New York, NY.
- NPCC, 2001. Northwest Power and Conservation Council. Technical Guidance for Subbasin Planners, Council Document 2001-20.
- NMFS (National Marine Fisheries Service). 1996. Making Endangered Species Act determinations of effect for individual or grouped actions at the watershed scale. The National Marine Fisheries Service Environmental and Technical Services Division, Habitat Conservation Branch, Seattle, WA.
- Pacific Coastal Salmon Recovery Fund, 2004. Final Data Dictionary.
- Peck, D. V., J. M. Lazorchak, and D. J. Klemm. 2001. Environmental monitoring and assessment program—surface waters: western pilot study field operations manual for wadeable streams. Draft Report. EPA/XXX/X-XX/XXX, U.S. Environmental Protection Agency, Washington, D.C. Web link: <http://www.epa.gov/emap/html/pubs/docs/groupdocs/surfwatr/field/ewwsm01.html>
- [Although this draft document states that it should not be cited or quoted, some of the material in the report is an important improvement to Lazorchak et al. (1998). By not citing the document, it may give the appearance that this document improves some of the methods outlined in the Lazorchak et al. report. To avoid this, PNAMP believes it is necessary to offer credit where credit is due.]*
- PNAMP, 2004. Recommendations for Coordinating State, Federal, and Tribal Watershed and Salmon Monitoring Programs in the Pacific Northwest.

- Reynolds, L., A. T. Herlihy, P. R. Kaufmann, S. V. Gregory, and R. M. Hughes. 2003. Electrofishing effort requirements for assessing species richness and biotic integrity in Western Oregon streams. *North American Journal of Fisheries Management* 23:450-461.
- Rosgen, D. 1996. Applied river morphology. Wildland Hydrology, Pagosa Springs, CO.
- Simonson, T., J. Lyons, and P. Kanehl. 1994. Quantifying fish habitat in streams: transect spacing, sample size, and a proposed framework. *North American Journal of Fisheries Management* 14:607-615.
- Spence, B. C., G. A. Lomnicky, R. M. Hughes, and R. P. Novitzki. 1996. An ecosystem approach to salmonid conservation. TR-4501-96-6057, Management Technology, Corvallis, OR.
- WMOC. 2002. The Washington Comprehensive Monitoring Strategy and Action Plan for watershed health and salmon recovery. Washington Monitoring Oversight Committee. Olympia, WA. Web link <http://www.iac.wa.gov/srfb/docs.htm>
- WSRFB (Washington Salmon Recovery Funding Board). 2003. Draft 5/16/2003 monitoring and evaluation strategy for habitat restoration and acquisition projects. Washington Salmon Recovery Funding Board, Olympia, WA. Web link: <http://www.iac.wa.gov/srfb/docs.htm>

Relevant References

Attached is an excerpted list of relevant references from Hillman et al., 2004. Since this contains many links to key documents and hosts a wealth of applicable citations, PNAMP has appended this to the guidance to help subbasin planners access this information.

- Anderson, R. O. and R. M. Neumann. 1996. Length, weight, and associated structural indices. Pages 447-482 *in*: B. R. Murphy and D. W. Willis, editors. *Fisheries techniques*, 2nd edition. American Fisheries Society, Bethesda, MD.
- Armantrout, N. B., compiler. 1998. Glossary of aquatic habitat inventory terminology. American Fisheries Society, Bethesda, MD.
- Bain, M. B. and N. J. Stevenson, editors. 1999. *Aquatic habitat assessment: common methods*. American Fisheries Society, Bethesda, MD.
- Bailey, R. G. 1978. Description of eco-regions of the United States. U.S. Forest Service, Intermountain Region, Ogden, UT.
- Bailey, R. G. 1998. Eco-regions map of North America: explanatory note. U.S. Forest Service, Miscellaneous Publication 1548, Washington, D.C.
- Bayley, P. B. 2002. A review of studies on responses of salmon and trout to habitat change, with potential for application in the Pacific Northwest. Report to the Washington State Independent Science Panel, Olympia, WA.
- Bevenger, G. S. and R. M. King. 1995. A pebble count procedure for assessing watershed cumulative effects. Research Paper RM-RP-319, USDA Forest Service, Rocky Mountain Forest and Range Experiment Station, Fort Collins, CO.
- Bilhimer, D., J. Carroll, S. O'Neal, and G. Pelletier. 2003. Draft quality assurance project plan: Wenatchee River temperature, dissolved oxygen, pH, and fecal coliform total maximum daily load year 2 technical study. Washington State Department of Ecology, Olympia, WA.

- Bisson, P. A. and D. R. Montgomery. 1996. Valley segments, stream reaches, and channel units. Pages 23-52 *in*: R. R. Hauer and G. A. Lamberti, editors. *Methods in stream ecology*. Academic Press, New York, NY.
- Borgerson, L. A. 1992. Scale analysis. Oregon Department of Fish and Wildlife, Fish Research Project F-144-R-4, Annual Progress Report, Portland, OR.
- Box, G. E. P. and G. M. Jenkins. 1976. *Time-series analysis: forecasting and control*. Holden-Day, San Francisco, CA.
- Browne, R. H. 2001. Using the sample range as a basis for calculating sample size in power calculations. *The American Statistician* 55:293-298.
- BURPTAC (Beneficial Use Reconnaissance Project Technical Advisory Committee). 1999. 1999 beneficial use reconnaissance project workplan for wadable streams. Idaho Division of Environmental Quality, Boise, ID. Web link: http://www.deq.state.id.us/water/surface_water/99_burp_workplan.pdf
- Cailliet, G. M., M. S. Love, and A. W. Ebeling. 1986. *Fishes, a field and laboratory manual on their structure, identification, and natural history*. Wadsworth Publishing Company, Belmont, CA.
- Clay, C. H. 1995. *Design of fishways and other fish facilities*. Second edition. Lewis Publishers, Boca Raton, Florida.
- Cohen, J. 1988. *Statistical power analysis for the behavioral sciences*. Lawrence-Erlbaum, Hillsdale, NJ.
- Cook, T. D. and D. T. Campbell. 1979. *Quasi-experimentation: design and analysis issues for field settings*. Houghton Mifflin Company, Boston, MA.
- Cupp, C. E. 1989a. Identifying spatial variability of stream characteristics through classification. Master's thesis. University of Washington, Seattle, WA.
- Cupp, C. E. 1989b. Valley segment type classification for forested lands of Washington. Washington State Timber/Fish/Wildlife Agreement, TFW-AM-89-001, Department of Natural Resources, Olympia, WA.
- Currens, K. P., H. W. Li, J. D. McIntyre, D. R. Montgomery, and D. W. Reiser. 2000. Recommendations for monitoring salmonid recovery in Washington State. Independent Science Panel Report 2000-2, Olympia, WA.
- Currens, K. P., H. W. Li, J. D. McIntyre, D. R. Montgomery, and D. W. Reiser. 2002. Responses of salmon and trout to habitat changes. Independent Science Panel Technical Memorandum 2002-2, Olympia, WA.
- Diaz-Ramos, S., D. L. Stevens, and A. R. Olsen. 1996. *EMAP statistical methods manual*. U.S. Environmental Protection Agency, EPA/620/R-96/XXX, Corvallis, OR.
- Dolloff, A., J. Kershner, and R. Thurow. 1996. Underwater observation. Pages 533-554 *in*: B. R. Murphy and D. W. Willis, editors. *Fisheries techniques*, 2nd edition. American Fisheries Society, Bethesda, MD.
- Frissell, C. A., W. J. Liss, C. E. Warren, and M. D. Hurley. 1986. A hierarchical framework for stream habitat classification; viewing streams in a watershed context. *Environmental Management* 10:199-214.
- Gordon, N. D., T. A. McMahon, and B. L. Finlayson. 1992. *Stream hydrology an introduction for ecologists*. John Wiley and Sons, New York, NY.
- Green, R. H. 1979. *Sampling design and statistical methods for environmental biologists*. John Wiley and Sons, Inc., New York, NY.

- Green, R. H. 1994. Aspects of power analysis in environmental monitoring. Pages 173-182 in: D. J. Fletcher and B. F. J. Manly, editors. *Statistics in ecology and environmental monitoring*. University of Otago Press, Dunedin.
- Hadley, R. F. and S. A. Schumm. 1961. Sediment sources and drainage basin characteristics in upper Cheyenne River basin. U.S. Geological Survey, Water-Supply Paper 1531-B, Reston, Virginia.
- Harrelson, C. C., C. L. Rawlins, and J. P. Potyondy. 1994. Stream channel reference sites: an illustrated guide to field technique. USDA Forest Service, Rocky Mountain Forest and Range Experiment Station, General Technical Report RM-245, Fort Collins, CO.
- Hillman, T. W., J. W. Mullan, and J. S. Griffith. 1992. Accuracy of underwater counts of juvenile Chinook salmon, coho salmon, and steelhead. *North American Journal of Fisheries Management* 12:598-603.
- Hillman, T. W. and D. W. Chapman. 1996. Comparison of underwater methods and electrofishing for estimating fish populations in the Upper Blackfoot River Basin. BioAnalysts, Inc. Report to the Seven-Up Pete Joint Venture, Lincoln, MT.
- Hillman, T. W. and M. D. Miller. 2002. Abundance and total numbers of Chinook salmon and trout in the Chiwawa River Basin, Washington. BioAnalysts, Inc. Report to Chelan County Public Utility District, Wenatchee, WA.
- Hughes, R. M., E. Rexstad, and C. E. Bond. 1987. The relationship of aquatic ecoregions, river basins and physiographic provinces to the ichthyogeographic regions of Oregon. *Copeia* 2:423-432.
- Hunt, C. B. 1967. *Physiography of the United States*. W. H. Freeman, San Francisco, CA.
- Hurlbert, S. J. 1984. Pseudoreplication and the design of ecological field experiments. *Ecological Monographs* 54:187-211.
- Idea Works, Inc. 1997. *Methodologists tool chest*. Version 1.2. Distributed by Scolaris, Sage Publications Software, Beverly Hills, CA.
- Kaufmann, P. R., P. Levine, E. G. Robinson, C. Seeliger, and D. V. Peck. 1999. Quantifying physical habitat in wadeable streams. EPA/620/R-99/003, U.S. Environmental Protection Agency, Washington, D.C. Web link: <http://www.epa.gov/emap/html/pubs/docs/groupdocs/surfwatr/field/phyhab.html>
- Lazorchak, J. M., D. J. Klemm, and D. V. Peck (editors). 1998. Environmental monitoring and assessment program—surface waters: field operations and methods for measuring the ecological condition of wadeable streams. EPA/620/R-94/004F, U.S. Environmental Protection Agency, Washington, D.C. Web link: http://www.epa.gov/emap/html/pubs/docs/groupdocs/surfwatr/field/ws_abs.html
- Lee, K. N. 1993. *Compass and gyroscope: integrating science and politics for the environment*. Island Press, Washington, D.C.
- Lipsey, M. W. 1990. *Design sensitivity: statistical power for experimental research*. Sage Publications, Beverly Hills, CA.
- MacDonald, L. H., A. W. Smart, and R. C. Wissmar. 1991. Monitoring guidelines to evaluate effects of forestry activities on streams in the Pacific Northwest and Alaska. U.S. Environmental Protection Agency, EPA/910/9-91-001, Seattle, WA.

- Manly, B. F. J. 1992. The design and analysis of research studies. Cambridge University Press, New York, NY.
- Mapstone, B. D. 1995. Scalable decision rules for environmental impact studies: effect size, Type I, and Type II errors. *Ecological Applications* 5:401-410.
- Meehan, W. R., editor. 1991. Influences of forest and range management on salmonid fishes and their habitats. American Fisheries Society Special Publication 19, Bethesda, MD.
- Meekin, T. K. 1967. Report on the 1966 Wells Dam chinook tagging study. Washington Department of Fisheries report to Douglas County Public Utility District, Contract Number 001-01-022-4201.
- Montgomery, D. R. and J. M. Buffington. 1993. Channel classification, prediction of channel response, and assessment of channel condition. Washington State Timber/Fish/Wildlife Agreement, TFW-SH10-93-002, Department of Natural Resources, Olympia, WA. Web site: http://www.nwifc.wa.gov/cmernoc/TFW_SH10_93_002.pdf
- Montgomery, D. R. and J. M. Buffington. 1997. Channel-reach morphology in mountain drainage basins. *Geological Society of American Bulletin* 109:596-611.
- Moore, K., K. Jones, and J. Dambacher. 1994. Methods for stream habitat surveys; aquatic inventory project. Oregon Department of Fish and Wildlife, Aquatic Inventories Project, Natural Production Program, Corvallis, OR. Web link: <http://oregonstate.edu/Dept/ODFW/freshwater/inventory/methods.html>
- Moore, K.M., K. F. Bierly and C. D. Pearson. 2002. Monitoring Strategy for the Oregon Plan for Salmon and Watersheds. Oregon Watershed Enhancement Board. Salem, Oregon.
- Mosey, T. R. and L. J. Murphy. 2002. Spring and summer Chinook spawning ground surveys on the Wenatchee River Basin, 2001. Chelan County Public Utility District, Wenatchee, WA.
- Mullan, J. W., K. R. Williams, G. Rhodus, T. W. Hillman, and J. D. McIntyre. 1992. Production and habitat of salmonids in Mid-Columbia River tributary streams. U.S. Fish and Wildlife Service Monograph I, Leavenworth, WA.
- Murdoch, A., K. Petersen, T. Miller, M. Tonseth, and T. Randolph. 2000. Freshwater production and emigration of juvenile spring chinook salmon from the Chiwawa River in 1998. Report No. SS99-05, Washington Department of Fish and Wildlife, Olympia, WA.
- Nawa, R. K., C. A. Frissell, and W. J. Liss. 1988. Life history and persistence of anadromous fish stocks in relation to stream habitats and watershed classification. Annual progress report to Oregon Department of Fish and Wildlife, Portland, OR.
- NCSS (Number Cruncher Statistical Systems). 2000. Pass 2000 power analysis and sample size for Windows. NCSS, Kaysville, UT.
- NRC (National Research Council). 1992. Restoration of aquatic ecosystems: science, technology, and public policy. National Academy Press, Washington, D.C.
- Omernik, J. M. 1987. Aquatic ecoregions of the conterminous United States. *Annals of the Association of American Geographers* 77:118-125.
- OPSW (Oregon Plan for Salmon and Watersheds). 1999. Water quality monitoring, technical guidebook. Version 2.0. Corvallis, OR. Web link: <http://www.oweb.state.or.us/publications/index.shtml>

- Overton, C. K., S. P. Wollrab, B. C. Roberts, and M. A. Radko. 1997. R1/R4 (Northern/Intermountain Regions) fish and fish habitat standard inventory procedures handbook. USDA Forest Service General Technical Report INT-GTR-346, Ogden, UT.
- Overton, W. S., D. White, and D. L. Stevens. 1990. Design report for EMAP environmental monitoring and assessment program. U.S. Environmental Protection Agency, EPA/600/3-91/053, Corvallis, OR.
- Parker, M. A. 2000. Fish passage – culvert inspection procedures. Watershed Restoration Technical Circular No. 11. Ministry of Environment, Lands and Parks and Ministry of Forest, British Columbia.
- Parker, R. A. and N. G. Berman. 2003. Sample size: more than calculations. *The American Statistician* 57:166-170.
- Parmenter, A. W., A. Hansen, R. E. Kennedy, W. Cohen, U. Langener, R. Lawrence, B. Maxwell, A. Gallant, and R. Aspinall. 2003. Land use and land cover in the greater Yellowstone ecosystem: 1975-1995. *Ecological Applications* 13:687-703.
- Paulsen, C., S. Katz, T. Hillman, A. Giorgi, C. Jordan, M. Newsom, and J. Geiselman, 2002. Guidelines for Action Effectiveness Research Proposals for FCRPS Offsite Mitigation Habitat Measures. Bonneville Power Administration, U.S. Bureau of Reclamation, U.S. Army Corps of Engineers, and NOAA Fisheries, Portland, OR. Web link:
<http://www.efw.bpa.gov/cgiin/FW/welcome.cgi?ViewMode=ExternalView>.
- Peterman, R. M. 1990. Statistical power analysis can improve fisheries research and management. *Canadian Journal of Fisheries and Aquatic Sciences* 47:2-15.
- Platts, W. S., W. F. Megahan, and G. W. Minshall. 1983. Methods for evaluating stream, riparian, and biotic conditions. USDA Forest Service General Technical Report INT-138, Ogden, UT.
- Platts, W. S. and twelve others. 1987. Methods for evaluating riparian habitats with applications to management. USDA Forest Service General Technical Report INT-221, Ogden, UT.
- Reeves, G. H., and nine others. 2001. Aquatic and riparian effectiveness monitoring plan for the Northwest Forest Plan. USDA Forest Service, Pacific Northwest Research Station, Corvallis, OR.
- Reynolds, J. B. 1996. Electrofishing. Pages 221-253 *in*: B. R. Murphy and D. W. Willis, editors. *Fisheries techniques*, 2nd edition. American Fisheries Society, Bethesda, MD.
- Roni, P., L. Weitkamp, and J. Scordino. 1999. Identification of essential fish habitat for salmon in the Pacific Northwest: initial efforts, information needs, and future direction. *American Fisheries Society Symposium* 22:93-107.
- Roper, B. B., J. L. Kershner, and R. C. Henderson. 2003. The value of using permanent sites when evaluating stream attributes at the reach scale. *Journal of Freshwater Ecology* 18:585-592.
- Royce, W. F. 1996. Introduction to the practice of fishery science. Revised edition. Academic Press, New York, NY.
- Scheaffer, R. L., W. Mendenhall, and L. Ott. 1990. Elementary survey sampling. Fourth edition. PWS-KENT Publishing Company, Boston, MA.

- Schuett-Hames, D., J. Ward, M. Fox, A. Pleus, and J. Light. 1994. Large woody debris survey module. Section 5 *in*: D. Schuett-Hames, A. Pleus, L. Bullchild, and S. Hall, editors. Ambient monitoring program manual. Timber-Fish-Wildlife TFW-AM9-94-001, Northwest Indian Fisheries Commission, Olympia, WA. Web link: <http://www.nwifc.wa.gov/TFW/documents.asp>
- Schuett-Hames, D., A. E. Pleus, E. Rashin, and J. Matthews. 1999a. Method manual for the stream temperature survey. Timber-Fish-Wildlife TFW-AM9-99-005, Northwest Indian Fisheries Commission, Olympia, WA. Web link: <http://www.nwifc.wa.gov/TFW/documents.asp>
- Schuett-Hames, D., R. Conrad, A. Pleus, and M. McHenry. 1999b. Method manual for the salmonid spawning gravel composition survey. Timber-Fish-Wildlife TFW-AM9-99-006, Northwest Indian Fisheries Commission, Olympia, WA. Web link: <http://www.nwifc.wa.gov/TFW/documents.asp>
- Skalski, J. R. and D. S. Robson. 1992. Techniques for wildlife investigations, design and analysis of capture data. Academic Press, New York, NY.
- Smith, E. P., D. R. Orvos, and J. Cairns. 1993. Impact assessment using the before-after-control-impact (BACI) model: concerns and comments. *Canadian Journal of Fisheries and Aquatic Sciences* 50:627-637.
- Smith, R. L. and T. M. Smith. 2001. Ecology and field biology. Sixth edition. Benjamin Cummings, New York, N.Y.
- Stevens, D. L. 1997. Variable density grid-based sampling designs for continuous spatial populations. *Environmetrics* 8:167-195.
- Stevens, D. L. 2002. Sampling design and statistical analysis methods for the integrated biological and physical monitoring of Oregon streams. Report No. OPSW-ODFW-2002-07, The Oregon Plan for Salmon and Watersheds, Oregon Department of Fish and Wildlife, Corvallis, OR.
- Stevens, D. L. and A. R. Olsen. 1999. Spatially restricted surveys over time for aquatic resources. *Journal of Agricultural, Biological, and Environmental Statistics* 4:415-428.
- Stevens, D. L. and N. S. Urquhart. 2000. Response designs and support regions in sampling continuous domains. *Environmetrics* 11:13-41.
- Stewart-Oaten, A., W. W. Murdoch, and K. R. Parker. 1986. Environmental impact assessment: "pseudoreplication" in time? *Ecology* 67:929-940.
- Stewart-Oaten, A., J. R. Bence, and C. W. Osenberg. 1992. Assessing effects of unreplicated perturbations: no simple solutions. *Ecology* 73:1396-1404.
- Strahler, A. N. 1952. Hypsometric (area-altitude) analysis of erosional topography. *Bulletin of the Geological Society of America* 63:1117-1142.
- Strange, R. J. 1996. Field examination of fishes. Pages 433-446 *in*: B. R. Murphy and D. W. Willis, editors. Fisheries techniques, 2nd edition. American Fisheries Society, Bethesda, MD.
- Underwood, A. J. 1994. Things environmental scientists (and statisticians) need to know to receive (and give) better statistical advice. Pages 33-61 *in*: D. J. Fletcher and B. F. J. Manly, editors. Statistics in ecology and environmental monitoring. University of Otago Press, Dunedin.

- USFWS (U.S. Fish and Wildlife Service). 2000. Biological Opinion, effects to listed species from operations of the Federal Columbia River Power System. U.S. Fish and Wildlife Service, Regions 1 and 6, Portland, OR.
- Van Deventer, J. S. and W. S. Platts. 1989. Microcomputer software system for generating population statistics from electrofishing data—user's guide for MicroFish 3.0. USDA Forest Service General Technical Report INT-254, Ogden, UT.
- WDFW (Washington Department of Fish and Wildlife). 2000. Fish passage barrier and surface water diversion screening assessment and prioritization manual. Washington Department of Fish and Wildlife Habitat Program, Environmental Restoration Division, Olympia, WA.
Web link: <http://www.wa.gov/wdfw/hab/engineer/fishbarr.htm>
- Wertz, W. A. and J. F. Arnold. 1972. Land systems inventory. U.S. Forest Service, Intermountain Region, Ogden, UT.
- WFC (World Forestry Center). 1998. Pilot study report, Umpqua land exchange project. World Forestry Center, Portland, OR. Web link:
<http://www.or.blm.gov/umpqua/documents.htm>
- WFPB (Washington Forest Practices Board). 1995. Washington forest practices board manual: standard methodology for conducting watershed analysis under Chapter 222-22 WAC. Version 3.0. Washington Forest Practices Board, Olympia, WA.
Web link: <http://www.dnr.wa.gov/forestpractices/watershedanalysis/>
- Whittier, T. R., R. M. Hughes, and D. P. Larsen. 1988. Correspondence between ecoregions and spatial patterns in stream ecosystems in Oregon. *Canadian Journal of Fisheries and Aquatic Sciences* 45:1264-1278.
- Wipfli, M. S. and D. P. Gregovich. 2002. Export of invertebrates and detritus from fishless headwater streams in southeastern Alaska: implications for downstream salmonid production. *Freshwater Biology* 47:957-969.
- Zaroban, D. W. 2000. Protocol for placement and retrieval of temperature data loggers in Idaho streams. Idaho Division of Environmental Quality, Boise, ID. Web link:
<http://www.deq.state.id.us/water/tlp.htm>